

**PCT**

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification 7 :  <b>C12N 15/52, 9/64, C07K 16/40, A61K 38/48, C12N 15/62, A01K 67/027, G01N 33/566, A61K 48/00</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 00/58479</b></p> <p>(43) International Publication Date: <b>5 October 2000 (05.10.00)</b></p>								
<p>(21) International Application Number: <b>PCT/US00/07755</b></p> <p>(22) International Filing Date: <b>23 March 2000 (23.03.00)</b></p> <p>(30) Priority Data:  <b>09/277,229      26 March 1999 (26.03.99)      US</b></p> <p>(71) Applicant: <b>AMGEN INC. [US/US]; One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).</b></p> <p>(72) Inventors: <b>CITRON, Martin; 1272 Wilder Street, Thousand Oaks, CA 91362 (US). VASSAR, Robert, James; 272 Via Colinas, Westlake Village, CA 91362 (US). BENNETT, Brian, Drake; 2706 Regina Avenue, Thousand Oaks, CA 91360 (US).</b></p> <p>(74) Agents: <b>ODRI, Steven, M. et al.; Amgen, Inc., One Amgen Center Drive, Thousand Oaks, CA 91320-1799 (US).</b></p>	<p>(81) Designated States: <b>AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</b></p> <p><b>Published</b>  <i>With international search report.          Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>									
<p>(54) Title: <b>BETA SECRETASE GENES AND POLYPEPTIDES</b></p> <div style="text-align: center;"> <table border="1" style="margin: 10px auto; border-collapse: collapse;"> <caption>Approximate data from bar chart</caption> <thead> <tr> <th>Condition</th> <th>Aβ42 (ng/ml)</th> </tr> </thead> <tbody> <tr> <td>vector</td> <td>~1.4</td> </tr> <tr> <td>β-sec</td> <td>~2.8</td> </tr> <tr> <td>β-sec D93A-HA</td> <td>~1.5</td> </tr> </tbody> </table> </div> <p style="text-align: center;"> <span style="display: inline-block; width: 10px; height: 10px; border: 1px solid black; background: repeating-linear-gradient(45deg, transparent, transparent 2px, black 2px, black 4px); margin-right: 5px;"></span> vector  <span style="display: inline-block; width: 10px; height: 10px; background: repeating-linear-gradient(-45deg, transparent, transparent 2px, black 2px, black 4px); margin-right: 5px;"></span> β-sec  <span style="display: inline-block; width: 10px; height: 10px; border: 1px solid black; margin-right: 5px;"></span> β-sec D93A-HA         </p>			Condition	Aβ42 (ng/ml)	vector	~1.4	β-sec	~2.8	β-sec D93A-HA	~1.5
Condition	Aβ42 (ng/ml)									
vector	~1.4									
β-sec	~2.8									
β-sec D93A-HA	~1.5									
<p>(57) Abstract</p> <p>Disclosed are novel genes encoding beta secretase polypeptides. Also disclosed are methods of making and using the polypeptides.</p>										

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LX	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

**Description**

5

10

15

20

25

30

35

40

45

50

55

- 1. -

## BETA SECRETASE GENES AND POLYPEPTIDES

Field of the Invention

The present invention relates to novel beta secretase polypeptides and nucleic acid molecules encoding the polypeptides. The invention also relates to vectors, host cells, antibodies and methods for producing beta secretase polypeptides.

Background of the Invention

Patients with Alzheimer's disease initially show short term memory loss. As the disease progresses, the patients become completely demented. The brains of Alzheimer's patients contain numerous amyloid plaques and neurofibrillary tangles (highly insoluble protein aggregates). Not surprisingly, such plaques and tangles are found primarily in those regions of the brain involved with memory and cognition.

The amyloid plaques are associated with dystrophic dendrites and axons, as well as activated microglia and reactive astrocytes. The major component of the plaques is a peptide referred to as A-beta (Selkoe, *Science*, 275:630-631 (1997)).

A-beta is ultimately generated by endoprotease cleavages of the polypeptide beta-amyloid precursor protein (beta-APP). Beta-APP is constitutively expressed in most cells of the body, but production of A-beta appears to be highest in brain cells. A-beta is typically secreted by the cells that produce it into the extracellular matrix where it makes its way into various bodily fluids. A schematic diagram of beta-APP processing is set forth in Figure 12.

- 2 -

There are three major isoforms of beta-APP polypeptide, and they are referred to as beta-APP695, beta-APP751 and beta-APP770. The number on each isoform refers to the number of amino acids it contains.

Beta-APP is synthesized as a membrane protein and spans the Golgi membrane. Normal non-pathogenic processing of beta-APP is believed to occur via a putative enzyme referred to as "alpha-secretase".

Alpha-secretase cleaves between amino acids 687 and 688 of beta-APP770 (or between amino acids 612 and 613 of beta-APP695), thereby generating a 687 amino acid soluble form of beta-APP770 (or a 612 amino acid soluble form of beta-APP695) referred to as "alpha-APPs". The remaining membrane-bound portion of beta-APP (amino acids 688-770 of the beta-APP770 isoform) is cleaved by the enzyme gamma-secretase. Gamma secretase cleaves between amino acids 711 and 712, and between amino acids 713 and 714, to generate the fragments 688-711, 688-713, and 712-770 and 714-770. The first two fragments are referred to as "p3" and are released from the membrane. Gamma secretase is also active on the two other isoforms of beta-APP.

In both non-diseased people and Alzheimer's disease patients, beta-APP is also processed via an alternative mechanism. Here, a heretofore uncharacterized enzyme referred to as "beta-secretase" cleaves full length beta-APP between amino acids 671 and 672 of beta-APP770, thereby generating a soluble fragment of amino acids 1-671, referred to as "beta-APPs" and a membrane bound form of amino acids 672-770. The membrane bound form is then acted on by gamma secretase which, as mentioned above, cleaves between amino acids 711 and 712, and between amino acids 713 and 714 to produce soluble fragments of amino acids 672-711 and 672-713; these fragments are referred to as

- 3 -

"A-beta 40", and "A-beta 42", respectively. In addition, gamma secretase activity generates membrane bound fragments of amino acids 712-770 and 714-770. A-beta 40 and A-beta 42 are secreted from the cells and accumulate in the brains of Alzheimer's patients to form the characteristic brain tissue plaques.

Recent evidence suggests that beta secretase can also cleave beta-APP between amino acids 683 and 684 of the 770 isoform (Gouras et al., *J. Neurochem.*, 71:1920-1925 [1998]) thereby ultimately generating a second form of A-beta spanning amino acids 684-711 and 684-713. The mechanisms that control this alternate cleavage are not known.

A few naturally occurring mutations of beta-APP have been identified. These include the so called "Swedish mutation" in which amino acids 670 and 671 of the 770 isoform are mutated, and the London mutation in which amino acid 717 of the 770 isoform is mutated. The Swedish mutation, which causes familial Alzheimer's disease, has been found to result in increased A-beta production *in vivo* (Citron et al., *Nature*, 360:672-674 [1992]; see also Citron et al., *Neuron*, 14:661-670 [1995]).

US Patent No. 5,744,346 purportedly describes a beta secretase molecule. No nucleic acid sequence or amino acid sequence for this molecule is presented.

European patent application EP O 855 444 A2 published 29 July 1998 describes an aspartic proteinase that purportedly plays a role in Alzheimer's disease. This aspartic proteinase has sequence similarity at both the nucleic acid and amino acid levels to the beta-secretase of the present invention.

European patent application EP O 848 062 A2, published 17 June 1998, describes an aspartic proteinase termed "ASP1" which has approximately 47 percent identity at the amino acid level to the prepro

- 4 -

form of the beta-secretase of the present invention as determined by a computer comparison using the GAP alignment program.

Accordingly, it is an object of the invention to identify nucleic acid molecules encoding beta secretase polypeptides. Such molecules have use as probes for diagnosis of Alzheimer's disease, and for identification of compounds that modulate the activity of beta secretase.

#### Summary of the Invention

The invention provides an isolated nucleic acid molecule selected from the group consisting of:

a) the nucleic acid molecule as set forth in any of SEQ ID NOS: 1, 2, and 3;

b) a nucleic acid molecule encoding the polypeptide of any of SEQ ID NOS: 4, 5, and 6;

c) biologically active fragments of SEQ ID NO:4;

d) an allelic variant or splice variant of any of (a) or (b);

e) a nucleic acid molecule of the DNA vector insert in ATCC Deposit No. 207158;

f) a nucleic acid molecule of the DNA vector insert in ATCC Deposit No. 207159;

g) a nucleic acid molecule encoding a polypeptide having one to fifty conservative amino acid substitutions as compared with the polypeptide of SEQ ID NO:4, wherein the polypeptide encoded by said nucleic acid molecule is biologically active; and

h) a nucleic acid molecule that is the complement of any of (a)-(g) above.

In other embodiments, the invention provides expression vectors, host cells, and methods of preparing recombinant beta secretase polypeptide.

5

- 5 -

10

The invention also provides an isolated polypeptide selected from the group consisting of:

15

a) the polypeptide of any of SEQ ID NOS: 4, 5, and 6;

5

b) a biologically active fragment of any of SEQ ID NOS. 4, 5, 6;

c) a biologically active polypeptide having one to fifty conservative amino acid changes as compared with the polypeptide of SEQ ID NO:4;

20

10

d) the polypeptide encoded by the DNA vector insert of ATCC Deposit Nos. 207158 and 207159; and

e) a polypeptide that is an allelic variant or splice variant of (a).

25

15

The invention further provides an isolated beta secretase polypeptide fragment of SEQ ID NO:4 selected from the group consisting of: amino acids 46-501; amino acids 46-460; amino acids 45-460; amino acids 1-460; amino acids 93-292; amino acids 93-293; amino acids 91-295; amino acids 90-295; amino acids 90-300; amino acids 280-310; amino acids 62-420; amino acids 1-420; amino acids 62-460; amino acids 90-460; amino acids 62-501; amino acids 62-460; amino acids 93-293; amino acids 90-293; amino acids 90-300; amino acids 62-420; amino acids 62-501; amino acids 1-420; amino acids 46-420; amino acids 62-420; amino acids 73-420; amino acids 83-420; amino acids 90-420; amino acids 62-417; amino acids 73-417; amino acids 83-417; amino acids 90-417; amino acids 62-410; amino acids 73-410; amino acids 83-410; amino acids 90-4104; amino acids 62-402; amino acids 73-402; amino acids 83-402; and amino acids 90-40.

30

20

35

25

40

45

The invention further provides an antibody or fragment thereof specifically binding to beta secretase polypeptide.

50

35

The invention yet further provides a fusion polypeptide comprising full length or truncated beta-

55



- 6 -

secretase polypeptide fused to a heterologous amino acid sequence such as the Fc portion of human IgG.

Description of the Figures

Figure 1A and 1B depict the cDNA sequence of human beta secretase (SEQ ID NO:1).

Figure 2A and 2B depict the cDNA sequence of mouse beta secretase (SEQ ID NO:2).

Figure 3A and 3B depict the cDNA sequence of rat beta secretase (SEQ ID NO:3).

Figure 4 depicts the putative amino acid sequence of human beta secretase polypeptide (SEQ ID NO:4).

Figure 5 depicts the putative amino acid sequence of mouse beta secretase polypeptide (SEQ ID NO:5).

Figure 6 depicts the putative amino acid sequence of rat beta secretase polypeptide (SEQ ID NO:6).

Figure 7 depicts a human multiple tissue Northern blot which was probed with a *Pst*I DNA fragment of human beta-secretase cDNA extending from nucleotide 318 to nucleotide 1090. Lane 1 is pancreas; Lane 2 is kidney; Lane 3 is skeletal muscle; Lane 4 is liver; Lane 5 is lung; Lane 6 is placenta; Lane 7 is brain; Lane 8 is heart; Lane 9 is peripheral blood leukocytes; Lane 10 is colon; Lane 11 is small intestine; Lane 12 is ovary; Lane 13 is spinal cord; Lane 14 is prostate; Lane 15 is thymus; Lane 16 is spleen. The lower panel shows a control hybridization using a DNA probe for

- 7 -

actin. RNA size markers (in kb) are indicated on the left.

Figure 3 depicts a Western blot of a SDS gel containing protein from either human 293 cells transfected with beta-secretase cDNA or human brain. Lane 1 contains high molecular weight standards; lane 2 contains immunoprecipitate of cell culture medium from human 293 cells transiently transfected with beta-secretase cDNA using preimmune serum; lane 3 contains immunoprecipitate of cell culture medium from human 293 cells transiently transfected with beta-secretase cDNA using beta-secretase antibody; lane 4 contains immunoprecipitate of homogenate from human Alzheimer's disease brain tissue using preimmune serum; lane 5 contains immunoprecipitate of homogenate from human Alzheimer's disease brain tissue using antibody against beta-secretase; lane 6 contains immunoprecipitate of homogenate from human control (non-Alzheimer's disease) brain tissue using preimmune serum; lane 7 contains immunoprecipitate of homogenate from human control (non-Alzheimer's disease) brain tissue using antibody against beta-secretase; lane 8 contains low molecular weight standards.

Figure 9A depicts a graph of ELISA assays to detect A-beta 42. APP expressing cells were transfected with either vector plasmid (negative control), the beta-secretase vector (positive control), or beta-secretase vector containing the D93A mutation.

Figure 9B depicts graph of ELISA assays to detect A-beta 40. APP expressing cells were transfected with either vector plasmid (negative control), the beta-secretase vector (positive control), or beta-secretase vector containing the D93A mutation.

- 8 -

Figure 10 depicts an SDS gel of purified human beta-secretase prepared as an Fc fusion. The beta-secretase-Fc fusion is indicated as "fusion" in Lane 2; "Fc" refers to an Fc fragment only. Molecular weight markers are indicated in Lane 1.

Figure 11 is a graph depicting the ability of human beta-secretase-Fc fusion to cleave various APP peptides. The X-axis is the amount of enzyme in the reaction mixture and the Y-axis is the amount of cleavage (denoted as pmol product formed) of each of three substrate peptides. "Sw" refers to the APP peptide with the Swedish mutation; "Wt" refers to wild type APP peptide; and "MV" refers to a mutated APP peptide.

Figure 12 is a schematic representation of the processing steps of beta-APP to generate A-beta. APP can undergo cleavage by alpha-secretase to form a secreted alpha-APP soluble fragment ("alpha-APPS") and a membrane bound fragment of about 10 kDa. The membrane bound fragment can then be cleaved by gamma-secretase to release a fragment referred to as "p3". Alternatively, APP can undergo beta-secretase cleavage to release a soluble fragment referred to as "beta-APPS" and a membrane bound fragment of about 12 kDa. The membrane bound fragment can then be cleaved by gamma-secretase to release A-beta.

#### Detailed Description of the Invention

The section headings herein are for organizational purposes only and are not to be construed as limiting the subject matter described therein.

- 9 -

Definitions

The term "beta secretase nucleic acid molecule" refers to a nucleic acid molecule comprising or consisting essentially of a nucleotide sequence as set forth in any of SEQ ID NOS: 1, 2, and 3, comprising or consisting essentially of a nucleotide sequence encoding any of the polypeptides as set forth in SEQ ID NOS: 4, 5, and 6, comprising or consisting essentially of a nucleotide sequence of the DNA insert in ATCC deposit number 207158 and ATCC deposit number 207159 (both deposited 11 March 1999), or nucleic acid molecules related thereto. Related nucleic acid molecules comprise or consist essentially of nucleotide sequences that are at least 70 percent identical to the nucleotide sequence as shown in any of SEQ ID NOS: 1, 2, and 3; or comprise or consist essentially of nucleotide sequences encoding polypeptides that are at least 70 percent identical to any of the polypeptides as set forth in SEQ ID NOS: 4, 5, and 6. The nucleotide sequences may be at least 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95 percent identical to any of the nucleotide sequences as shown in SEQ ID NOS: 1, 2, and 3, or the nucleotide sequences that encode polypeptides that are about 75 percent, or about 80 percent, or about 85 percent, or about 90 percent, or about 95 percent identical to any of the polypeptide sequences as set forth in SEQ ID NOS 4, 5, and 6. Related nucleic acid molecules also include fragments of the above beta secretase nucleic acid molecules which are at least about 10 contiguous nucleotides, or about 15, or about 20, or about 25, or about 50, or about 75, or about 100, or greater than about 100 contiguous nucleotides. Related nucleic acid molecules also include fragments of the above beta secretase nucleic acid molecules

- 10 -

5 which encode polypeptides of at least about 25 amino  
10 acid residues, or about 50, or about 75, or about 100,  
or greater than about 100 amino acid residues. Related  
beta secretase nucleic acid molecules include those  
5 molecules which comprise nucleotide sequences which  
15 hybridize under moderate or highly stringent conditions  
as defined herein with any of the above nucleic acid  
molecules. In preferred embodiments, the related  
nucleic acid molecules comprise sequences which  
20 hybridize under moderate or highly stringent conditions  
with the sequence as shown in SEQ ID NO:1, or with a  
molecule encoding a polypeptide, which polypeptide  
comprises the sequence as shown in SEQ ID NO:4, or with  
25 a nucleic acid fragment as defined above, or with a  
nucleic acid fragment encoding a polypeptide as defined  
above. It is also understood that related nucleic acid  
molecules include allelic or splice variants of any of  
30 the above nucleic acids, and include sequences which  
are complementary to any of the above nucleotide  
20 sequences.

The term "isolated nucleic acid molecule"  
refers to a nucleic acid molecule of the invention that  
35 is free from at least one contaminating nucleic acid  
molecule with which it is naturally associated, and  
25 preferably substantially free from any other  
contaminating mammalian nucleic acid molecules which  
would interfere with its use in protein production or  
40 its therapeutic or diagnostic use.

The term "allelic variant" refers to one of  
30 several possible naturally occurring alternate forms of  
a gene occupying a given locus on a chromosome of an  
45 organism.

The term "splice variant" refers to a  
nucleic acid molecule, usually RNA, which is generated  
35 by alternative processing of intron sequences in an RNA  
transcript.  
50  
55

- 11 -

The term "expression vector" refers to a vector which is suitable for propagation in a host cell and contains nucleic acid sequences which direct and/or control the expression of inserted heterologous nucleic acid sequences. Expression includes, but is not limited to, processes such as transcription, translation, and RNA splicing, if introns are present.

The term "high stringency conditions" refers to those conditions that (1) employ low ionic strength reagents and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate/0.1% NaDodSO<sub>4</sub> (SDS) at 50°C. or (2) employ during hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0.1%. Alternatively, Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 may be used with 750 mM NaCl, 75 mM sodium citrate at 42°C. Another example is the use of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS.

The term "moderate stringency conditions" refers to conditions which generally include the use of a washing solution and hybridization conditions (e.g., temperature, ionic strength, and percent SDS) less stringent than described above. An example of moderately stringent conditions are conditions such as overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 µl/ml

- 12 -

denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength etc. as necessary to accommodate factors such as probe length and the like.

Where oligonucleotide probes are used to screen cDNA or genomic libraries, one of the following two high stringency solutions may be used. The first of these is 6 X SSC with 0.05 percent sodium pyrophosphate at a temperature of 35°C-62°C, depending on the length of the oligonucleotide probe. For example, 14 base pair probes are washed at 35-40°C, 17 base pair probes are washed at 45-50°C, 20 base pair probes are washed at 52-57°C, and 23 base pair probes are washed at 57-63°C. The temperature can be increased 2-3°C where the background non-specific binding appears high. A second high stringency solution utilizes tetramethylammonium chloride (TMAC) for washing oligonucleotide probes. One stringent washing solution is 3 M TMAC, 50 mM Tris-HCl, pH 8.0, and 0.2 percent SDS. The washing temperature using this solution is a function of the length of the probe. For example, a 17 base pair probe is washed at about 45-50°C.

The term "beta secretase polypeptide" refers to a polypeptide comprising the amino acid sequence of any of SEQ ID NOS: 4, 5, and 6, and related polypeptides described herein. Related polypeptides includes allelic variants, splice variants, fragments, derivatives, substitution, deletion, and insertion variants, fusion polypeptides, and orthologs. Beta secretase polypeptides may be processed polypeptides, i.e., not containing an endogenous or exogenous signal or leader sequence as defined herein, and may or may

- 13 -

not have an amino terminal methionine residue,  
depending on the method by which they are prepared.

The term "mature beta-secretase polypeptide"  
refers to a polypeptide of any of SEQ ID NOS: 4, 5 and  
6 and related polypeptides described herein, in which  
the leader sequence and the propeptide have been  
removed. Mature human beta-secretase is amino acids  
46-501 of SEQ ID NO:4. For human beta-secretase, the  
leader peptide is amino acids 1-21 of SEQ ID NO:4, and  
the propeptide is amino acids 22-45 of SEQ ID NO:4.

The term "beta secretase polypeptide  
fragment" refers to a peptide or polypeptide that  
comprises less than the full length amino acid sequence  
of an beta secretase polypeptide as set forth in any of  
SEQ ID NOS: 4, 5, and 6. Such a fragment may arise,  
for example, from a truncation at the amino terminus, a  
truncation at the carboxy terminus, and/or an internal  
deletion of a residue(s) from the amino acid sequence.  
Naturally occurring beta secretase fragments may result  
from alternative RNA splicing, from in vivo processing  
such as removal of the leader peptide and propeptide,  
and/or from protease activity.

The term "beta secretase polypeptide  
variants" refers to beta secretase polypeptides  
comprising amino acid sequences which contain one or  
more amino acid sequence substitutions, deletions,  
and/or additions as compared to the beta secretase  
polypeptide amino acid sequence set forth in any of SEQ  
ID NOS: 4, 5, and 6 or fragments thereof. Variants may  
be naturally occurring or artificially constructed.  
Such beta secretase polypeptide variants may be  
prepared from the corresponding nucleic acid molecules  
encoding said variants, which have a DNA sequence that  
varies accordingly from the DNA sequences for wild type  
beta secretase polypeptides as set forth in any of SEQ  
ID NOS: 4, 5, and 6.



- 14 -

5  
10 The term "beta secretase fusion polypeptide" refers to a fusion of beta secretase polypeptide, fragment, variant and/or derivative thereof, with a heterologous peptide or polypeptide.

15 The term "beta secretase polypeptide derivatives" refers to beta secretase polypeptides, variants, or fragments thereof, that have been chemically modified, as for example, by covalent attachment of one or more water soluble polymers,  
20 N-linked or O-linked carbohydrates, sugars, phosphates, and/or other such molecules. The derivatives are modified in a manner that is different from naturally occurring beta secretase, either in the type and/or location of the molecules attached to the polypeptide  
25 Derivatives further include the deletion of one or more chemical groups naturally attached to the beta secretase polypeptide.

30 The terms "biologically active beta secretase polypeptides", "biologically active beta secretase polypeptide fragments", "biologically active beta secretase polypeptide variants", and "biologically active beta secretase polypeptide derivatives" refer to  
35 beta secretase polypeptides having at least one activity characteristic of a beta secretase polypeptide, such as the ability to cleave the APP Swedish mutation peptide EVKMDAEF (SEQ ID NO:18)  
40 between the methionine and aspartic acid residues.

45 The term "isolated polypeptide" refers to a polypeptide of the invention that is free from at least one contaminating polypeptide that is found in its natural environment, and preferably substantially free from any other contaminating mammalian polypeptides which would interfere with its therapeutic or diagnostic use.

50 The term "ortholog" refers to a polypeptide that corresponds to a polypeptide identified from a

55

- 15 -

different species. For example, mouse and human beta secretase polypeptides are considered orthologs.

The terms "effective amount" and "therapeutically effective amount" refer to the amount of a beta secretase polypeptide that is useful or necessary to support an observable level one or more biological activities of the beta secretase polypeptides as set forth above.

10 Relatedness of Nucleic Acid Molecules and/or Polypeptides

The term "identity", as known in the art, refers to a relationship between the sequences of two or more polypeptide molecules or two or more nucleic acid molecules, as determined by comparing the sequences. In the art, "identity" also means the degree of sequence relatedness between polypeptide or nucleic acid molecule sequences, as the case may be, as determined by the match between strings of nucleotide or amino acid sequences. "Identity" measures the percent of identical matches between two or more sequences with gap alignments addressed by a particular mathematical model or computer programs (i.e., "algorithms").

The term "similarity" is a related concept, but in contrast to "identity", refers to a measure of similarity which includes both identical matches and conservative substitution matches. Since conservative substitutions apply to polypeptides and not nucleic acid molecules, similarity only deals with polypeptide sequence comparisons. If two polypeptide sequences have, for example, 10/20 identical amino acids, and the remainder are all non-conservative substitutions, then the percent identity and similarity would both be 50%. If in the same example, there are 5 more positions where there are conservative substitutions, then the

- 16 -

percent identity remains 50%, but the per cent  
similarity would be 75% (15/20). Therefore, in cases  
where there are conservative substitutions, the degree  
of similarity between two polypeptide sequences will be  
higher than the percent identity between those two  
sequences.

The term "conservative amino acid  
substitution" refers to a substitution of a native  
amino acid residue with a nonnative residue such that  
there is little or no effect on the polarity or charge  
of the amino acid residue at that position. For  
example, a conservative substitution results from the  
replacement of a non-polar residue in a polypeptide  
with any other non-polar residue. Furthermore, any  
native residue in the polypeptide may also be  
substituted with alanine, as has been previously  
described for "alanine scanning mutagenesis". General  
classes of amino acids useful for conservative amino  
acid substitutions are set forth in Table I.

17 -

Table I

## Conservative Amino Acid Substitutions

Basic:	arginine
	lysine
	histidine
Acidic:	glutamic acid.
	aspartic acid
Uncharged Polar:	glutamine
	asparagine
	serine
	threonine
	tyrosine
Non-Polar:	phenylalanine
	tryptophan
	cysteine
	glycine
	alanine
	valine
	proline
	methionine
	leucine
	norleucine
	isoleucine

Conservative modifications to the amino acid sequence (and the corresponding modifications to the encoding nucleotides) are expected to produce beta secretase having functional and chemical characteristics similar to those of naturally occurring beta secretase.

Conservative amino acid substitutions also encompass non-naturally occurring amino acid residues

- 18 -

which are typically incorporated by chemical peptide synthesis rather than by synthesis in biological systems. Such conservative amino acids include, for example, the "homolog" of each amino acid, where the homolog is an amino acid with a methylene group (CH<sub>2</sub>) inserted into the side chain at the beta position of that side chain.

Beta-secretase polypeptides of the present invention may have 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, 45, 50, or up to 55 conservative amino acid changes as compared with the beta-secretase polypeptide of SEQ ID NO:4. Such molecules may also have chemical modifications as described herein for beta-secretase variants.

In contrast, non-conservative substitutions of beta secretase may be generated by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the molecular backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Non-conservative substitutions may involve the exchange of a member of one of the amino acid classes of Table I for a member from another class. Such substituted residues may be introduced into regions of the human beta secretase molecule that are homologous with non-human beta secretase, or into the non-homologous regions of the molecule.

Identity and similarity of related nucleic acid molecules and polypeptides can be readily calculated by known methods, including but not limited to those described in *Computational Molecular Biology*, Lesk, A.M., ed., Oxford University Press, New York, 1988; *Biocomputing: Informatics and Genome Projects*, Smith, D.W., ed., Academic Press, New York, 1993;

- 19 -

Computer Analysis of Sequence Data. Part 1. Griffin, A.M., and Griffin, H.G., eds., Humana Press, New Jersey, 1994; *Sequence Analysis in Molecular Biology*, von Heinje, G., Academic Press, 1987; and *Sequence Analysis Primer*, Gribskov, M. and Devereux, J., eds., M. Stockton Press, New York, 1991; and Carillo, H., and Lipman, D., *SIAM J. Applied Math.*, 48:1073 (1988).

Preferred methods to determine identity and/or similarity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include, but are not limited to, the GCG program package, including GAP (Devereux, et al., *Nucleic Acids Research* 12:387 [1984]; Genetics Computer Group, University of Wisconsin, Madison, WI), BLASTP, BLASTN, and FASTA (Atschul et al., *J. Molec. Biol.* 215:403-410 [1990]). The BLAST X program is publicly available from the National Center for Biotechnology Information (NCBI) and other sources (*BLAST Manual*, Altschul et al., NCB NLM NIH Bethesda, MD 20894; Altschul et al., *J. Mol. Biol.* 215:403-410 [1990]). The well known Smith Waterman algorithm may also be used to determine identity.

By way of example, using the computer algorithm GAP (Genetics Computer Group, University of Wisconsin, Madison, WI), two polypeptides for which the percent sequence identity is to be determined are aligned for optimal matching of their respective amino acids (the "matched span", as determined by the algorithm). A gap opening penalty (which is calculated as 3 X the average diagonal; the "average diagonal" is the average of the diagonal of the comparison matrix being used; the "diagonal" is the score or number

- 20 -

assigned to each perfect amino acid match by the particular comparison matrix) and a gap extension penalty (which is usually 1/10 times the gap opening penalty), as well as a comparison matrix such as PAM 250 or BLOSUM 62 are used in conjunction with the algorithm. A standard comparison matrix (see Dayhoff et al., in: Atlas of Protein Sequence and Structure, vol. 5, supp. 3 [1978] for the PAM250 comparison matrix; see Henikoff et al., *Proc. Natl. Acad. Sci. USA*, 89:10915-10919 [1992] for the BLOSUM 62 comparison matrix) is also used by the algorithm.

Preferred parameters for polypeptide sequence comparison include the following:

- Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)
- Comparison matrix: BLOSUM 62 from Henikoff and Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915-10919 (1992)
- Gap Penalty: 12
- Gap Length Penalty: 4
- Threshold of Similarity: 0

The GAP program is useful with the above parameters. The aforementioned parameters are the default parameters for polypeptide comparisons (along with no penalty for end gaps) using the GAP algorithm.

Preferred parameters for nucleic acid molecule sequence comparison include the following:

- Algorithm: Needleman and Wunsch, *J. Mol. Biol.* 48:443-453 (1970)
- Comparison matrix: matches = +10, mismatch = 0

- 21 -

- Gap Penalty: 50
- Gap Length Penalty: 3

The GAP program is also useful with the above parameters. The aforementioned parameters are the default parameters for nucleic acid molecule comparisons.

Other exemplary algorithms, gap opening penalties, gap extension penalties, comparison matrices, thresholds of similarity, etc. may be used by those of skill in the art, including those set forth in the Program Manual, Wisconsin Package, Version 9, September, 1997. The particular choices to be made will depend on the specific comparison to be made, such as DNA to DNA, protein to protein, protein to DNA; and additionally, whether the comparison is between given pairs of sequences (in which case GAP or BestFit are generally preferred) or between one sequence and a large database of sequences (in which case FASTA or BLASTA are preferred).

#### Nucleic Acid Molecules

Recombinant DNA methods used herein are generally those set forth in Sambrook et al. (*Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY [1989]) and/or Ausubel et al., eds., (*Current Protocols in Molecular Biology*, Green Publishers Inc. and Wiley and Sons, NY [1994]).

The invention provides for nucleic acid molecules as described herein and methods for obtaining the molecules. A gene or cDNA encoding a beta secretase polypeptide or fragment thereof may be obtained by hybridization screening of a genomic or cDNA library, or by PCR amplification. Probes or



- 22 -

primers useful for screening a library by hybridization can be generated based on sequence information for other known genes or gene fragments from the same or a related family of genes, such as, for example, conserved motifs. In addition, where a gene encoding beta secretase polypeptide has been identified from one species, all or a portion of that gene may be used as a probe to identify corresponding genes from other species (orthologs) or related genes from the same species (homologs). The probes or primers may be used to screen cDNA libraries from various tissue sources believed to express the beta secretase gene. In addition, part or all of a nucleic acid molecule having any of the sequences as set forth in SEQ ID NOS: 1, 2, and 3 may be used to screen a genomic library to identify and isolate a gene encoding beta secretase. Typically, conditions of moderate or high stringency will be employed for screening to minimize the number of false positives obtained from the screen.

Nucleic acid molecules encoding beta secretase polypeptides may also be identified by expression cloning which employs detection of positive clones based upon a property of the expressed protein. Typically, nucleic acid libraries are screened by binding of an antibody or other binding partner (e.g., receptor or ligand) to cloned proteins which are expressed and displayed on the host cell surface. The antibody or binding partner is modified with a detectable label to identify those cells expressing the desired clone.

Another means of preparing a nucleic acid molecule encoding a beta secretase polypeptide or fragment thereof is chemical synthesis using methods well known to the skilled artisan such as those described by Engels et al. (*Angew. Chem. Intl. Ed.*, 28:716-734 [1989]). These methods include, *inter alia*,

- 23 -

the phosphotriester, phosphoramidite, and H-phosphonate methods for nucleic acid synthesis. A preferred method for such chemical synthesis is polymer-supported synthesis using standard phosphoramidite chemistry.

Typically, the DNA encoding the beta secretase polypeptide will be several hundred nucleotides in length. Nucleic acids larger than about 100 nucleotides can be synthesized as several fragments using these methods. The fragments can then be ligated together to form the full length beta secretase polypeptide. Usually, the DNA fragment encoding the amino terminus of the polypeptide will have an ATG, which encodes a methionine residue. This methionine may or may not be present on the mature form of the beta secretase polypeptide, depending on whether the polypeptide produced in the host cell is designed to be secreted from that cell.

In some cases, it may be desirable to prepare nucleic acid molecules encoding beta secretase polypeptide variants. Nucleic acid molecules encoding variants may be produced using site directed mutagenesis, PCR amplification, or other appropriate methods, where the primer(s) have the desired point mutations (see Sambrook et al., *supra*, and Ausubel et al., *supra*, for descriptions of mutagenesis techniques). Chemical synthesis using methods described by Engels et al., *supra*, may also be used to prepare such variants. Other methods known to the skilled artisan may be used as well.

In one embodiment, nucleic acid molecule variants contain codons which have been altered for optimal expression of an beta secretase polypeptide in a given host cell. Particular codon alterations will depend upon the beta secretase polypeptide(s) and host cell(s) selected for expression. Such "codon optimization" can be carried out by a variety of

- 24 -

5 methods, for example, by selecting codons which are  
10 preferred for use in highly expressed genes in a given  
host cell. Computer algorithms which incorporate codon  
frequency tables such as "Ecohigh. Cod" for codon  
5 preference of highly expressed bacterial genes may be  
used and are provided by the University of Wisconsin  
15 Package Version 9.0, Genetics Computer Group, Madison,  
WI. Other useful codon frequency tables include  
"Celegans\_high.cod", "Celegans\_low.cod",  
10 "Drosophila\_high.cod", "Human\_high.cod",  
20 "Maize\_high.cod", and "Yeast\_high.cod".

In other embodiments, nucleic acid molecules  
encode beta secretase variants with conservative amino  
acid substitutions as defined above, beta secretase  
25 variants comprising an addition and/or a deletion of  
one or more N-linked or O-linked glycosylation sites,  
or beta secretase polypeptide fragments as described  
above. In addition, nucleic acid molecules may encode  
30 any combination of beta secretase variants, fragments,  
20 and fusion polypeptides described herein.

Preferred nucleic acid molecule fragments  
encode the following fragments of SEQ ID NO:4: amino  
35 acids 45-501; amino acids 46-501; amino acids 46-460;  
amino acids 45-460; amino acids 1-460; amino acids 93-  
25 292; amino acids 93-293; amino acids 91-295; amino  
acids 90-295; amino acids 90-300; amino acids 62-420;  
40 amino acids 1-420; amino acids 62-460; amino acids 90-  
460; amino acids 62-501; amino acids 62-460; amino  
acids 90-293; amino acids 90-300; amino acids 1-420;  
30 amino acids 46-420; amino acids 62-420; amino acids 73-  
420; amino acids 83-420; amino acids 90-420; amino  
45 acids 62-417; amino acids 73-417; amino acids 83-417;  
amino acids 90-417; amino acids 62-410; amino acids 73-  
410; amino acids 83-410; amino acids 90-410; amino  
35 acids 62-402; amino acids 73-402; amino acids 83-402;  
50 and amino acids 90-402.

55

- 25 -

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

Additionally preferred nucleic acid molecule fragments include nucleic acid molecules encoding each of the foregoing fragments joined to a nucleic acid molecule encoding a peptide or polypeptide such as, for example, the Fc portion of human IgG.

#### Vectors and Host Cells

A nucleic acid molecule encoding a beta secretase polypeptide can be inserted into an appropriate expression vector in order to generate beta secretase polypeptides. The vector is typically selected to be functional in the particular host cell employed (i.e., the vector is compatible with the host cell machinery such that amplification of the gene and/or expression of the gene can occur). A nucleic acid molecule encoding a beta secretase polypeptide may be amplified/expressed in prokaryotic, yeast, insect (baculovirus systems) and/or eukaryotic host cells. Selection of the host cell will depend in part on whether an beta secretase polypeptide is to be post-translationally modified (e.g., glycosylated and/or phosphorylated). If so, yeast, insect, or mammalian host cells are preferable.

Typically, expression vectors used in any of the host cells will contain sequences for plasmid maintenance and for cloning and expression of exogenous nucleotide sequences. Such sequences, collectively referred to as "flanking sequences" will typically include one or more of the following nucleic acid molecules: a promoter, one or more enhancer sequences, an origin of replication, a transcriptional termination sequence, a complete intron sequence containing a donor and acceptor splice site, a leader sequence for secretion, a ribosome binding site, a polyadenylation sequence, a polylinker region for inserting the nucleic acid encoding the polypeptide to be expressed, and a

- 26 -

selectable marker element. Each of these sequences is discussed below.

Optionally, the vector may contain a "tag" sequence, i.e., an oligonucleotide molecule located at the 5' or 3' end of the beta secretase polypeptide coding sequence; the oligonucleotide molecule encodes polyHis (such as hexaHis), or other "tag" such as FLAG, HA (hemagglutinin Influenza virus) or myc for which commercially available antibodies exist. This tag is typically fused to the polypeptide upon expression of the polypeptide, and can serve as a means for affinity purification of the beta secretase polypeptide from the host cell. Affinity purification can be accomplished, for example, by column chromatography using antibodies against the tag as an affinity matrix. Optionally, the tag can subsequently be removed from the purified beta secretase polypeptide by various means such as using one or more selected peptidases for cleavage.

Flanking sequences may be homologous (i.e., from the same species and/or strain as the host cell), heterologous (i.e., from a species other than the host cell species or strain), hybrids (i.e., a combination of flanking sequences from more than one source), or synthetic, or native sequences which normally function to regulate beta secretase expression. As such, the source of flanking sequences may be any prokaryotic or eukaryotic organism, any vertebrate or invertebrate organism, or any plant, provided that the flanking sequence is(are) functional in, and can be activated by, the host cell machinery.

The flanking sequences useful in the vectors of this invention may be obtained by any of several methods well known in the art. Typically, flanking sequences useful herein other than the beta secretase gene flanking sequences will have been previously identified by mapping and/or by restriction

5

- 27 -

10

15

endonuclease digestion and can thus be isolated from the proper tissue source using the appropriate restriction endonucleases. In some cases, the full nucleotide sequence of one or more flanking sequence may be known. Here, the flanking sequence may be synthesized using the methods described above for nucleic acid synthesis or cloning.

20

Where all or only a portion of the flanking sequence is known, it may be obtained using PCR and/or by screening a genomic library with suitable oligonucleotide and/or flanking sequence fragments from the same or another species.

25

30

35

Where the flanking sequence is not known, a fragment of DNA containing a flanking sequence may be isolated from a larger piece of DNA that may contain, for example, a coding sequence or even another gene or genes. Isolation may be accomplished by restriction endonuclease digestion to produce the proper DNA fragment followed by isolation using agarose gel purification, Qiagen® column chromatography, or other method known to the skilled artisan. Selection of suitable enzymes to accomplish this purpose will be readily apparent to one of ordinary skill in the art.

40

45

An origin of replication is typically a part of prokaryotic expression vectors purchased commercially, and aids in the amplification of the vector in a host cell. Amplification of the vector to a certain copy number can, in some cases, be important for optimal expression of the beta secretase polypeptide. If the vector of choice does not contain an origin of replication site, one may be chemically synthesized based on a known sequence, and ligated into the vector.

50

The origin of replication from the plasmid pBR322 (Product No. 303-3S, New England Biolabs, Beverly, MA) is suitable for most Gram-negative

55

- 28 -

bacteria. Various origin of replication elements (e.g., SV40, polyoma, adenovirus, vesicular stomatitis virus (VSV) or papillomaviruses such as HPV or BPV) are useful for cloning vectors in mammalian cells.

Generally, the origin of replication component is not needed for mammalian expression vectors (for example, the SV40 origin is often used only because it contains the early promoter).

A transcription termination sequence is typically located 3' of the end of a polypeptide coding regions and serves to terminate transcription. Usually, a transcription termination sequence in prokaryotic cells is a G-C rich fragment followed by a poly T sequence. While the sequence is easily cloned from a library or even purchased commercially as part of a vector, it can also be readily synthesized using methods for nucleic acid synthesis such as those described above.

A selectable marker gene element encodes a protein necessary for the survival and growth of a host cell grown in a selective culture medium. Typical selection marker genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, tetracycline, or kanamycin for prokaryotic host cells, (b) complement auxotrophic deficiencies of the cell; or (c) supply critical nutrients not available from complex media. Preferred selectable markers are the kanamycin resistance gene, the ampicillin resistance gene, and the tetracycline resistance gene.

Other selection genes may be used to amplify the gene which will be expressed. Amplification is the process wherein genes which are in greater demand for the production of a protein critical for growth are reiterated in tandem within the chromosomes of successive generations of recombinant cells. Examples

- 29 -

of suitable selectable markers for mammalian cells include dihydrofolate reductase (DHFR) and thymidine kinase. The mammalian cell transformants are placed under selection pressure which only the transformants are uniquely adapted to survive by virtue of the marker present in the vector. Selection pressure is imposed by culturing the transformed cells under conditions in which the concentration of selection agent in the medium is successively changed, thereby leading to amplification of both the selection gene and the DNA that encodes beta secretase. As a result, increased quantities of beta secretase are synthesized from the amplified DNA.

A ribosome binding site is usually necessary for translation initiation of mRNA and is characterized by a Shine-Dalgarno sequence (prokaryotes) or a Kozak sequence (eukaryotes). The element is typically located 3' to the promoter and 5' to the coding sequence of the beta secretase polypeptide to be expressed. The Shine-Dalgarno sequence is varied but is typically a polypurine (i.e., having a high A-G content). Many Shine-Dalgarno sequences have been identified, each of which can be readily synthesized using methods set forth above and used in a prokaryotic vector.

A leader, or signal, sequence may be used to direct an beta secretase polypeptide out of the host cell. Typically, the signal sequence is positioned in the coding region of the beta secretase nucleic acid molecule, or directly at the 5' end of the beta secretase polypeptide coding region.

The signal sequence may be a component of the vector, or it may be a part of beta secretase DNA that is inserted into the vector. The native beta secretase DNA encodes a signal sequence at the amino terminus of the protein that is cleaved during post-translational



- 30 -

processing of the molecule to form the mature beta secretase protein product.

Included within the scope of this invention are beta secretase nucleic acid molecules with the native signal sequence as well as beta secretase nucleic acid molecules wherein the native signal sequence is deleted and replaced with a heterologous signal sequence. The heterologous signal sequence selected should be one that is recognized and processed, i.e., cleaved by a signal peptidase, by the host cell. For expression of beta secretase in prokaryotic host cells, the native signal sequence is typically replaced by a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, or heat-stable enterotoxin II leaders. For beta secretase expression in yeast host cells, the native beta secretase signal sequence may be substituted by any of the yeast invertase, alpha factor, or acid phosphatase signal sequences. For beta secretase expression in mammalian host cells, the native beta secretase signal sequence is satisfactory, although other mammalian signal sequences may be suitable.

In many cases, transcription of a nucleic acid molecule is increased by the presence of one or more introns in the vector; this is particularly true where a polypeptide is produced in eukaryotic host cells, especially mammalian host cells. The introns used may be naturally occurring within the beta secretase gene, especially where the gene used is a full length genomic sequence or a fragment thereof. Where the intron is not naturally occurring within the gene (as for most cDNAs), the intron(s) may be obtained from another source. The position of the intron with respect to flanking sequences and the beta secretase gene is generally important, as the intron must be

- 31 -

transcribed to be effective. Thus, when an beta secretase cDNA molecule is being expressed, the preferred position for the intron is 3' to the transcription start site, and 5' to the polyA transcription termination sequence. Preferably, the intron or introns will be located on one side or the other (i.e., 5' or 3') of the cDNA such that it does not interrupt the coding sequence. Any intron from any source, including any viral, prokaryotic and eukaryotic (plant or animal) organisms, may be used to practice this invention, provided that it is compatible with the host cell(s) into which it is inserted. Also included herein are synthetic introns. Optionally, more than one intron may be used in the vector.

The expression and cloning vectors of the present invention will typically contain a promoter that is recognized by the host organism and operably linked to the molecule encoding the beta secretase polypeptide. Promoters are untranslated nucleic acid molecules located upstream(5') to the start codon of a structural gene (generally within about 100 to 1000 bp) that control the transcription and translation of a particular molecule, such as the beta secretase gene. Promoters are conventionally grouped into one of two classes, inducible promoters and constitutive promoters. Inducible promoters initiate increased levels of transcription of the structural gene in response to some change in culture conditions, such as the presence or absence of a nutrient or a change in temperature.

A large number of promoters that are functional in various host cells are well known and readily available. These promoters can be operably linked to the DNA encoding beta secretase by removing the native promoter from the beta secretase gene (via restriction enzyme digestion) and inserting the desired

- 32 -

promoter into the vector. While the native beta secretase promoter sequence may be used to direct amplification and/or expression of the beta secretase gene in mammalian cells, a heterologous promoter is preferred if it permits greater transcription and higher yields of beta secretase polypeptide as compared to the native beta secretase promoter, and if it is compatible with the host cell system that has been selected for use.

Promoters suitable for use with prokaryotic hosts include, without limitation, the beta-lactamase and lactose promoter systems; alkaline phosphatase, a tryptophan (trp) promoter system; and hybrid promoters such as the tac promoter. Other known bacterial promoters are also suitable. Their sequences have been published, thereby enabling one skilled in the art to ligate them to the desired DNA sequence(s), using linkers or adapters as needed to supply any required restriction sites.

Suitable promoter sequences for use with yeast host cells are also well known in the art. Yeast enhancers are advantageously used with yeast promoters. Suitable promoters for use with mammalian host cells are well known and include those obtained from the genomes of viruses such as polyoma virus, fowlpox virus, adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and most preferably Simian Virus 40 (SV40).

Suitable mammalian host cell promoters include heterologous mammalian promoters, e.g., heat-shock promoters and the actin promoter.

Additional promoters which may be of interest in controlling beta secretase expression include, but are not limited to: the SV40 early promoter region; the CMV promoter; the promoter contained in the 3' long

- 33 -

terminal repeat of Rous sarcoma virus; the herpes  
thymidine kinase promoter; the regulatory sequences of  
the metallothionine gene; prokaryotic expression  
vectors such as the beta-lactamase promoter; or the tac  
promoter. Also of interest are the following animal  
transcriptional control regions, which exhibit tissue  
specificity and have been utilized in transgenic  
animals: the insulin gene control region which is  
active in pancreatic beta cells; the immunoglobulin  
gene control region which is active in lymphoid cells;  
the mouse mammary tumor virus control region which is  
active in testicular, breast, lymphoid and mast cells;  
albumin gene control region which is active in liver;  
the alphafetoprotein gene control region which is  
active in liver; the alpha 1-antitrypsin gene control  
region which is active in the liver; the beta-globin  
gene control region which is active in myeloid cells;  
the myelin basic protein gene control region which is  
active in oligodendrocyte cells in the brain; the  
myosin light chain-2 gene control region which is  
active in skeletal muscle; and the gonadotropic  
releasing hormone gene control region which is active  
in the hypothalamus.

Another optional component of the vector used  
for beta secretase polypeptide expression is the  
enhancer element. This nucleic acid molecule may be  
inserted into the vector to increase the transcription  
in higher eukaryotic host cells of a genomic DNA or  
cDNA molecule encoding beta secretase polypeptide.  
Enhancers are usually about 10-300 nucleotides in  
length and act on the promoter to increase its  
transcription. Several enhancer sequences available  
from mammalian genes are known (e.g., globin, elastase,  
albumin, alpha-feto-protein and insulin). Typically,  
however, an enhancer from a virus will be used. The  
SV40 enhancer, the cytomegalovirus early promoter

- 34 -

enhancer, the polyoma enhancer, and adenovirus enhancers are exemplary enhancers useful for the activation of eukaryotic promoters. While an enhancer may be inserted into the vector either 5' or 3' to the beta secretase gene or cDNA, it is typically located at a site 5' from the promoter.

Expression vectors of the invention may be constructed from a starting vector such as a commercially available vector. Such vectors may or may not contain all of the desired flanking sequences. Where one or more of the flanking sequences set forth above are not already present in the vector to be used, they may be individually obtained and ligated into the vector. Methods used for obtaining each of the flanking sequences are well known to one skilled in the art.

Preferred vectors for practicing this invention are those which are compatible with bacterial, insect, and mammalian host cells. Such vectors include, inter alia, pCRII, pCR3, and pcDNA3.1 (Invitrogen Company, San Diego, CA), pBSII (Stratagene Company, La Jolla, CA), pET15b (Novagen, Madison, WI), pGEX (Pharmacia Biotech, Piscataway, NJ), pEGFP-N2 (Clontech, Palo Alto, CA), pETL (BlueBacII; Invitrogen), and pFastBacDual (Gibco/BRL, Grand Island, NY).

Additional possible vectors include, but are not limited to, cosmids, plasmids or modified viruses, but the vector system must be compatible with the selected host cell. Such vectors include, but are not limited to plasmids such as Bluescript<sup>®</sup> plasmid derivatives (a high copy number ColE1-based phagemid, Stratagene Cloning Systems Inc., La Jolla CA), PCR cloning plasmids designed for cloning Taq-amplified PCR products (e.g., TOPO<sup>™</sup> TA Cloning<sup>®</sup> Kit, PCR2.1<sup>®</sup> plasmid

- 35 -

derivatives, Invitrogen, Carlsbad, CA), and mammalian, yeast or virus vectors such as a baculovirus expression system (pBacPAK plasmid derivatives, Clontech, Palo Alto, CA). The recombinant molecules can be introduced into host cells via transformation, transfection, infection, electroporation, or other known techniques.

After the vector has been constructed and a nucleic acid molecule encoding an beta secretase polypeptide has been inserted into the proper site of the vector, the completed vector may be inserted into a suitable host cell for amplification and/or polypeptide expression.

Host cells may be prokaryotic host cells (such as *E. coli*) or eukaryotic host cells (such as a yeast cell, an insect cell, or a vertebrate cell). The host cell, when cultured under appropriate conditions, synthesizes an beta secretase polypeptide which can subsequently be collected from the culture medium (if the host cell secretes it into the medium) or directly from the host cell producing it (if it is not secreted). Selection of an appropriate host cell will depend upon various factors, such as desired expression levels, polypeptide modifications that are necessary for activity, such as glycosylation or phosphorylation, and ease of folding into a biologically active molecule.

Suitable host cells or cell lines may be mammalian cells, such as Chinese hamster ovary cells (CHO), human embryonic kidney (HEK) 293 or 293T cells, or 3T3 cells. The selection of suitable mammalian host cells and methods for transformation, culture, amplification, screening and product production and purification are known in the art. Other suitable mammalian cell lines, are the monkey COS-1 and COS-7 cell lines, and the CV-1 cell line. Further exemplary mammalian host cells include primate cell lines and

- 36 -

rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from in vitro culture of primary tissue, as well as primary explants, are also suitable. Candidate cells may be genotypically deficient in the selection gene, or may contain a dominantly acting selection gene. Other suitable mammalian cell lines include but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, 3T3 lines derived from Swiss, Balb-c or NIH mice, BHK or HaK hamster cell lines. Each of these cell lines is known by and available to those skilled in the art of protein expression.

Similarly useful as host cells suitable for the present invention are bacterial cells. For example, the various strains of *E. coli* (e.g., HB101, DH5a, DH10, and MC1061 are well-known as host cells in the field of biotechnology. Various strains of *B. subtilis*, *Pseudomonas spp.*, other *Bacillus spp.*, *Streptomyces spp.*, and the like may also be employed in this method.

Many strains of yeast cells known to those skilled in the art are also available as host cells for expression of the polypeptides of the present invention. Preferred yeast cells include, for example, *Saccharomyces cerevisiae*.

Additionally, where desired, insect cell systems may be utilized in the methods of the present invention. Such systems are described for example in Kitts et al. (*Biotechniques*, 14:810-817 [1993]), Lucklow (*Curr. Opin. Biotechnol.*, 4:564-572 [1993]) and Lucklow et al. (*J. Virol.*, 67:4566-4579 [1993]). Preferred insect cells are Sf-9 and Hi5 (Invitrogen, Carlsbad, CA).

Transformation or transfection of an expression vector for an beta secretase polypeptide into a selected host cell may be accomplished by well

- 37 -

known methods including methods such as calcium chloride, electroporation, microinjection, lipofection or the DEAE-dextran method. The method selected will in part be a function of the type of host cell to be used. These methods and other suitable methods are well known to the skilled artisan, and are set forth, for example, in Sambrook et al., *supra*.

#### Polypeptide Production

Host cells comprising a beta secretase expression vector may be cultured using standard media well known to the skilled artisan. The media will usually contain all nutrients necessary for the growth and survival of the cells. Suitable media for culturing *E. coli* cells are for example, Luria Broth (LB) and/or Terrific Broth (TB). Suitable media for culturing eukaryotic cells are RPMI 1640, MEM, DMEM, all of which may be supplemented with serum and/or growth factors as required by the particular cell line being cultured. A suitable medium for insect cultures is Grace's medium supplemented with yeastolate, lactalbumin hydrolysate, and/or fetal calf serum as necessary.

Typically, an antibiotic or other compound useful for selective growth of transfected or transformed cells is added as a supplement to the media. The compound to be used will be dictated by the selectable marker element present on the plasmid with which the host cell was transformed. For example, where the selectable marker element is kanamycin resistance, the compound added to the culture medium will be kanamycin.

The amount of a beta secretase polypeptide produced by a host cell can be evaluated using standard methods known in the art. Such methods include, without limitation, Western blot analysis,



- 38 -

SDS-polyacrylamide gel electrophoresis, non-denaturing gel electrophoresis, HPLC separation, immunoprecipitation, and/or activity assays such as DNA binding gel shift assays.

If a beta secretase polypeptide has been designed to be secreted from the host cells, the majority of polypeptide may be found in the cell culture medium. If however, the beta secretase polypeptide is not secreted from the host cells, it will be present in the cytoplasm and/or the nucleus (for eukaryotic host cells) or in the cytosol (for gram negative bacteria host cells).

For a beta secretase polypeptide situated in the host cell cytoplasm and/or nucleus, the host cells are typically first disrupted mechanically or with detergent to release the intra-cellular contents into a buffered solution. Beta secretase polypeptide can then be isolated from this solution.

Purification of a beta secretase polypeptide from solution can be accomplished using a variety of techniques. If the polypeptide has been synthesized such that it contains a tag such as hexahistidine or other small peptide such as FLAG (Eastman Kodak Co., New Haven, CT) or myc (Invitrogen, Carlsbad, CA) at either its carboxyl or amino terminus, it may essentially be purified in a one-step process by passing the solution through an affinity column where the column matrix has a high affinity for the tag or for the polypeptide directly (i.e., a monoclonal antibody specifically recognizing beta secretase polypeptide). For example, polyhistidine binds with great affinity and specificity to nickel, thus an affinity column of nickel (such as the Qiagen® nickel columns) can be used for purification of beta secretase polypeptide/polyHis. (See for example, Ausubel et al.,

- 39 -

eds., *Current Protocols in Molecular Biology*, Section  
19.11.8, John Wiley & Sons, New York [1993]).

Where a beta secretase polypeptide is prepared without a tag attached and no antibodies are available, other well known procedures for purification can be used. Such procedures include, without limitation, ion exchange chromatography, molecular sieve chromatography, HPLC, native gel electrophoresis in combination with gel elution, and preparative isoelectric focusing ("Isoprime" machine/technique, Hoefer Scientific). In some cases, two or more of these techniques may be combined to achieve increased purity.

If a beta secretase polypeptide is produced intracellularly, the intracellular material (including inclusion bodies for gram-negative bacteria) can be extracted from the host cell using any standard technique known to the skilled artisan. For example, the host cells can be lysed to release the contents of the periplasm/cytoplasm by French press, homogenization, and/or sonication followed by centrifugation.

If a beta secretase polypeptide has formed inclusion bodies in the cytosol, the inclusion bodies can often bind to the inner and/or outer cellular membranes and thus will be found primarily in the pellet material after centrifugation. The pellet material can then be treated at pH extremes or with chaotropic agent such as a detergent, guanidine, guanidine derivatives, urea, or urea derivatives in the presence of a reducing agent such as dithiothreitol at alkaline pH or tris carboxyethyl phosphine at acid pH to release, break apart, and solubilize the inclusion bodies. The beta secretase polypeptide in its now soluble form can then be analyzed using gel electrophoresis, immunoprecipitation or the like. If

- 40 -

it is desired to isolate the beta secretase polypeptide, isolation may be accomplished using standard methods such as those set forth below and in Marston et al. (*Meth. Enz.*, 182:264-275 [1990]).

In some cases, a beta secretase polypeptide may not be biologically active upon isolation. Various methods for "refolding" or converting the polypeptide to its tertiary structure and generating disulfide linkages, can be used to restore biological activity. Such methods include exposing the solubilized polypeptide to a pH usually above 7 and in the presence of a particular concentration of a chaotropic agent. In most cases the refolding/oxidation solution will also contain a reducing agent or a reducing agent plus its oxidized form in a specific ratio to generate a particular redox potential allowing for disulfide shuffling to occur in the formation of cysteine bridge(s) of the polypeptide. Some commonly used redox couples include cysteine/cystamine, glutathione (GSH)/dithiobis GSH, cupric chloride, dithiothreitol (DTT)/dithiane DTT, 2-mercaptoethanol (BME)/dithio-b(BME). In many instances a cosolvent is necessary to increase the efficiency of the refolding and the more common reagents used for this purpose include glycerol, polyethylene glycol of various molecular weights, arginine and the like.

If inclusion bodies are not formed to a significant degree upon expression of an beta secretase polypeptide, the polypeptide will be found primarily in the supernatant after centrifugation of the cell homogenate and may be further isolated from the supernatant using methods such as those set forth below.

In situations where it is preferable to partially or completely purify an beta secretase polypeptide such that it is partially or substantially

5

- 41 -

10

15

20

25

30

35

40

45

50

55

free of contaminants, standard methods known to the one skilled in the art may be used. Such methods include, without limitation, separation by electrophoresis followed by electroelution, various types of chromatography (affinity, immunoaffinity, molecular sieve, and/or ion exchange), and/or high pressure liquid chromatography. In some cases, it may be preferable to use more than one of these methods for complete purification.

Beta secretase polypeptides, fragments, and/or derivatives thereof may also be prepared by chemical synthesis methods using techniques known in the art such as those set forth by Merrifield et al., (*J. Am. Chem. Soc.*, 85:2149 (1963)), Houghten et al. (*Proc Natl Acad. Sci. USA*, 82:5132 (1985)), and Stewart and Young (*Solid Phase Peptide Synthesis*, Pierce Chemical Co., Rockford, IL (1984)). Using these methods, beta secretase fragments of up to about 75 amino acids in length can be prepared.

Typically, beta-secretase fragments and variants are synthesized from readily available starting materials. Synthesis is usually conducted from carboxy to amino terminus. During synthesis, the alpha-amine of the amino acid to be added is protected by a urethane such as Boc, Cbz, Fmoc, or Alloc (see Greene et al., *Protective Groups in Organic Synthesis*, 2d. ed., John Wiley and Sons (1991) for a list of protective groups) while the free carboxyl is activated with an activating reagent which is usually a carbodiimide such as DCC (Dicyclohexyl carbodiimide), EDC (1-(3-dimethylaminopropyl)-3-ethylcarbodiimide), or DIC (diisopropylcarbodiimide). A preferred protective group is Fmoc. The activating reagent can optionally be used in the presence of a catalyst such as Hobt (N-Hydroxybenzotriazole) Hoat (7-aza-N-

- 42 -

hydroxybenzotriazole), Hosu, or Dmap (Dimethylaminopyridine). After the peptide is completely synthesized, the side chain protecting groups may be removed using methods set forth in the above cited references. Such methods include, without limitation, hydrogenation in the presence of a catalyst such as palladium, platinum, or rhodium; treatment with sodium in liquid ammonia, hydrochloric, hydrofluoric, hydrobromic, formic, trifluoromethanesulfonic, or trifluoroacetic acid; secondary amines; fluoride ion; trimethylsilyl halides such as bromide and iodide; or alkali.

The above described methods may be accomplished manually or using an automated peptide synthesizer such as an Applied Biosystems model 430, 430A, A431, or A433, using programming modules as defined by the manufacturer's manuals.

To generate beta secretase fragments or variants using chemical synthesis that are longer than about 75 amino acids, a technique known as chemical ligation can be used in which peptides can be ligated together. This technique is described by Baca et al. (*J. Amer. Chem. Soc.*, 117:1881-1887 [1995]) and by Schrolver et al. (*Int. J. Peptide Protein Res.*, 40:180-193 [1992]).

Chemically synthesized beta-secretase polypeptides, fragments, and variants may be oxidized to permit the formation of disulfide bridges using standard methods set forth in the above cited references.

The beta-secretase polypeptides or fragments thereof are expected to have biological activity comparable to beta-secretase polypeptides or fragments thereof produced recombinantly and thus may be used interchangeably with recombinant or beta-secretase peptide.

- 43 -

Another means of obtaining beta secretase polypeptide is via purification from biological samples such as source tissues and/or fluids in which the beta secretase polypeptide is naturally found. Such purification can be conducted using methods for protein purification as described above. The presence of the beta secretase polypeptide during purification may be monitored using, for example, an antibody prepared against recombinantly produced beta secretase polypeptide or peptide fragments thereof.

#### Polypeptides

Polypeptides of the invention include isolated beta secretase polypeptides and polypeptides related thereto including fragments, variants, fusion polypeptides, and derivatives as defined hereinabove.

Beta secretase fragments of the invention may result from truncations at the amino terminus (with or without a leader sequence), truncations at the carboxy terminus, and/or deletions internal to the polypeptide. In preferred embodiments, truncations and/or deletions comprise about 10 amino acids, or about 20 amino acid, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or more than about 100 amino acids. The polypeptide fragments so produced will comprise about 25 contiguous amino acids, or about 50 amino acids, or about 75 amino acids, or about 100 amino acids, or about 150 amino acids, or about 200 amino acids. Such beta secretase polypeptides fragments may optionally comprise an amino terminal methionine residue.

Preferred beta secretase polypeptide fragments of SEQ ID NO: 4 include: amino acids 45-501; amino acids 46-501; amino acids 46-460; amino acids 45-460; amino acids 1-460; amino acids 93-292; amino acids 93-293; amino acids 91-295; amino acids 90-295; amino

- 44 -

acids 90-300; amino acids 62-420; amino acids 1-420;  
amino acids 62-460; amino acids 90-460; amino acids 62-  
501; amino acids 62-460; amino acids 90-293; amino  
acids 90-300; amino acids 1-420; amino acids 46-420;  
amino acids 62-420; amino acids 73-420; amino acids 83-  
420; amino acids 90-420; amino acids 62-417; amino  
acids 73-417; amino acids 83-417; amino acids 90-417;  
amino acids 62-410; amino acids 73-410; amino acids 83-  
410; amino acids 90-410; amino acids 62-402; amino  
acids 73-402; amino acids 83-402; and amino acids 90-  
402.

Additionally preferred fragments include each  
of the foregoing prepared as a fusion peptide with a  
second peptide or polypeptide such as, for example, the  
Fc portion of human IgG.

Beta secretase polypeptide variants of the  
invention include one or more amino acid substitutions,  
additions and/or deletions as compared to any of SEQ ID  
NOS: 4, 5, and 6. In preferred embodiments, the  
variants have from 1 to 3, or from 1 to 5, or from 1 to  
10, or from 1 to 15, or from 1 to 20, or from 1 to 25,  
or from 1 to 50, or from 1 to 75, or from 1 to 100, or  
more than 100 amino acid substitutions, insertions,  
additions and/or deletions, wherein the substitutions  
may be conservative, as defined above, or non-  
conservative or any combination thereof. The variants  
may have additions of amino acid residues either at the  
carboxy terminus or at the amino terminus (with or  
without a leader sequence).

Preferred beta secretase polypeptide variants  
include glycosylation variants wherein the number  
and/or type of glycosylation sites has been altered  
compared to native beta secretase polypeptide. In one  
embodiment, beta secretase variants comprise a greater  
or a lesser number of N-linked glycosylation sites. An  
N-linked glycosylation site is characterized by the

- 45 -

sequence: Asn-X-Ser or Thr, where the amino acid residue designated as X may be any type of amino acid except proline. Substitution(s) of amino acid residues to create this sequence provides a potential new site for addition of an N-linked carbohydrate chain.

Alternatively, substitutions to eliminate this sequence will remove an existing N-linked carbohydrate chain.

Also provided is a rearrangement of N-linked carbohydrate chains wherein one or more N-linked glycosylation sites (typically those that are naturally occurring) are eliminated and one or more new N-linked sites are created.

Beta secretase fusion polypeptides of the invention comprise beta secretase polypeptides, fragments, variants, or derivatives fused to a heterologous peptide or protein. Heterologous peptides and proteins include, but are not limited to, an epitope to allow for detection and/or isolation of an beta secretase fusion polypeptide, a transmembrane receptor protein or a portion thereof, such as an extracellular domain, or a transmembrane and intracellular domain, a ligand or a portion thereof which binds to a transmembrane receptor protein, an enzyme or portion thereof which is catalytically active, a protein or peptide which promotes oligomerization, such as leucine zipper domain, and a protein or peptide which increase stability, such as an immunoglobulin constant region. An beta secretase polypeptide may be fused to itself or to a fragment, variant, or derivative thereof. Fusions may be made either at the amino terminus or at the carboxy terminus of an beta secretase polypeptide, and may be direct with no linker or adapter molecule or may be through a linker or adapter molecule, such as one or more amino acid residues up to about 20 amino acids residues, or up to about 50 amino acid residues. A linker or



- 46 -

adapter molecule may also be designed with a cleavage site for a DNA restriction endonuclease or for a protease to allow for separation of the fused moieties.

In a preferred embodiment, a beta secretase polypeptide, fragment, variant and/or derivative is fused to an Fc region of human IgG. In one example, a human IgG hinge, CH2 and CH3 region may be fused at either the N-terminus or C-terminus of the beta secretase polypeptides using methods known to the skilled artisan. In another example, a portion of a hinge regions and CH2 and CH3 regions may be fused. The beta secretase Fc-fusion polypeptide so produced may be purified by use of a Protein A affinity column. In addition, peptides and proteins fused to an Fc region have been found to exhibit a substantially greater half-life in vivo than the unfused counterpart. Also, a fusion to an Fc region allows for dimerization/multimerization of the fusion polypeptide.

Beta secretase polypeptide derivatives are included in the scope of the present invention. Such derivatives are chemically modified beta secretase polypeptide compositions in which beta secretase polypeptide is linked to a polymer. The polymer selected is typically water soluble so that the protein to which it is attached does not precipitate in an aqueous environment, such as a physiological environment. The polymer selected is usually modified to have a single reactive group, such as an active ester for acylation or an aldehyde for alkylation, so that the degree of polymerization may be controlled as provided for in the present methods. The polymer may be of any molecular weight, and may be branched or unbranched. Included within the scope of beta secretase polypeptide polymers is a mixture of polymers. Preferably, for therapeutic use of the end-

- 47 -

product preparation, the polymer will be pharmaceutically acceptable.

The water soluble polymer or mixture thereof may be selected from the group consisting of, for example, polyethylene glycol (PEG), monomethoxy-polyethylene glycol, dextran, cellulose, or other carbohydrate based polymers, poly-(N-vinyl pyrrolidone) polyethylene glycol, propylene glycol homopolymers, a polypropylene oxide/ethylene oxide co-polymer, polyoxyethylated polyols (e.g., glycerol) and polyvinyl alcohol.

For the acylation reactions, the polymer(s) selected should have a single reactive ester group. For reductive alkylation, the polymer(s) selected should have a single reactive aldehyde group. A preferred reactive aldehyde is polyethylene glycol propionaldehyde, which is water stable, or mono C1-C10 alkoxy or aryloxy derivatives thereof (see U.S. Patent No. 5,252,714).

Pegylation of beta secretase polypeptides may be carried out by any of the pegylation reactions known in the art, as described for example in the following references: *Focus on Growth Factors 3*: 4-10 (1992); EP 0 154 316; and EP 0 401 384. Preferably, the pegylation is carried out via an acylation reaction or an alkylation reaction with a reactive polyethylene glycol molecule (or an analogous reactive water-soluble polymer) as described below.

A particularly preferred water-soluble polymer for use herein is polyethylene glycol, abbreviated PEG. As used herein, polyethylene glycol is meant to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono-(C1-C10) alkoxy- or aryloxy-polyethylene glycol.

In general, chemical derivatization may be performed under any suitable conditions used to react a

- 48 -

biologically active substance with an activated polymer molecule. Methods for preparing pegylated beta secretase polypeptides will generally comprise the steps of (a) reacting the polypeptide with polyethylene glycol (such as a reactive ester or aldehyde derivative of PEG) under conditions whereby beta secretase polypeptide becomes attached to one or more PEG groups, and (b) obtaining the reaction product(s). In general, the optimal reaction conditions for the acylation reactions will be determined based on known parameters and the desired result. For example, the larger the ratio of PEG: protein, the greater the percentage of poly-pegylated product.

In a preferred embodiment, the beta secretase polypeptide derivative will have a single PEG moiety at the N terminus.

Generally, conditions which may be alleviated or modulated by administration of the present beta secretase polypeptide derivative include those described herein for beta secretase polypeptides. However, the beta secretase polypeptide derivative disclosed herein may have additional activities, enhanced or reduced biological activity, or other characteristics, such as increased or decreased half-life, as compared to the non-derivatized molecules.

#### Antibodies

Beta secretase polypeptides, fragments, variants and derivatives may be used to prepare antibodies using methods known in the art. Thus, antibodies and antibody fragments that bind beta secretase polypeptides are within the scope of the present invention. Antibodies may be polyclonal, monoclonal, recombinant, chimeric, humanized, fully human, single chain and/or bispecific.

- 49 -

Polyclonal antibodies directed toward a beta secretase polypeptide generally are raised in animals (rabbits or mice) by multiple subcutaneous or intraperitoneal injections of beta secretase in combination with an adjuvant. It may be useful to conjugate a beta secretase polypeptide, or a variant, fragment or derivative thereof to a carrier protein that is immunogenic in the species to be immunized, such as keyhole limpet heccyanin, serum, albumin, bovine thyroglobulin, or soybean trypsin inhibitor. Also, aggregating agents such as alum are used to enhance the immune response. After immunization, the animals can be bled and the serum is assayed for anti-beta secretase antibody titer.

Monoclonal antibodies directed toward beta secretase polypeptide can be produced using any method which provides for the production of antibody molecules by continuous cell lines in culture. Examples of suitable methods for preparing monoclonal antibodies include hybridoma method of Kohler et al., *Nature* 256: 495-497 (1975), and the human B-cell hybridoma method, Kozbor, *J. Immunol.* 133: 3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, pp. 51-63 (Marcel Dekker, Inc., New York, 1987).

Also provided by the invention are hybridoma cell lines which produce monoclonal antibodies reactive with beta secretase polypeptides.

Monoclonal antibodies of the invention may be modified for use as therapeutics. One embodiment is a "chimeric" antibody in which a portion of the heavy and/or light chain is identical with or homologous to corresponding sequence in antibodies derived from a particular species or belonging to a particular antibody class or subclass, while the remainder of the chain(s) is identical with or homologous to

- 50 -

corresponding sequence in antibodies derived from another species or belonging to another antibody class or subclass, as well as fragments of such antibodies, so long as they exhibit the desired biological activity (see U.S. Patent No. 4,816,567; and Morrison et al., *Proc. Natl. Acad. Sci.* 81, 6851-6855 [1985]).

Also included in the scope of the present invention are monoclonal antibodies that are "humanized". Methods for humanizing non-human antibodies are well known in the art. Humanization can be performed following methods known in the art (see Jones et al., *Nature* 321: 522-525 [1986]; Riechmann et al., *Nature*, 332: 323-327 [1988]; Verhoeven et al., *Science* 239: 1534-1536 [1988]), by substituting rodent complementarily-determining regions (CDRs) for the corresponding regions of a human antibody.

Also encompassed by the invention are fully human antibodies which bind beta secretase polypeptides, fragments, variants and/or derivatives. Such antibodies can be produced by immunization with an beta secretase antigen (optionally conjugated to a carrier) of transgenic animals (e.g., mice) that are capable of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. See, for example, Jakobovits et al., *Proc. Natl. Acad. Sci.* 90: 2551-2555 (1993); Jakobovits et al., *Nature* 362: 255-258 (1993); Bruggermann et al., *Year in Immuno.* 7:33 (1993). Human antibodies can also be produced in phage-display libraries (see Hoogenboom et al., *J. Mol. Biol.* 227:381 [1991]; Marks et al., *J. Mol. Biol.* 222:581 [1991]).

For diagnostic applications, anti-beta secretase antibodies typically will be labeled with a detectable moiety. The detectable moiety can be any one which is capable of producing, either directly or indirectly, a detectable signal. For example, the

- 51 -

detectable moiety may be a radioisotope, such as  $^3\text{H}$ ,  $^{14}\text{C}$ ,  $^{32}\text{P}$ ,  $^{35}\text{S}$ , or  $^{125}\text{I}$ , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin; or an enzyme, such as alkaline phosphatase,  $\beta$ -galactosidase or horseradish peroxidase.

The anti-beta secretase antibodies of the invention may be employed in any known assay method, such as competitive binding assays, direct and indirect sandwich assays, and immunoprecipitation assays (see Sola, *Monoclonal Antibodies: A Manual of Techniques*, pp. 147-158 [CRC Press, Inc., 1987]) for detection and quantitation of beta secretase polypeptides. The antibodies will bind beta secretase polypeptides with an affinity which is appropriate for the assay method being employed.

Competitive binding assays rely on the ability of a labeled standard (e.g., an beta secretase polypeptide, or an immunologically reactive portion thereof) to compete with the test sample analyte (an beta secretase polypeptide) for binding with a limited amount of anti beta secretase antibody. The amount of an beta secretase polypeptide in the test sample is inversely proportional to the amount of standard that becomes bound to the antibodies. To facilitate determining the amount of standard that becomes bound, the antibodies generally are insolubilized before or after the competition, so that the standard and analyte that are bound to the antibodies may conveniently be separated from the standard and analyte which remain unbound.

Sandwich assays involve the use of two antibodies, each capable of binding to a different immunogenic portion, or epitope, of the protein to be detected and/or quantitated. In a sandwich assay, the test sample analyte is typically bound by a first

- 52 -

antibody which is immobilized on a solid support, and thereafter a second antibody binds to the analyte, thus forming an insoluble three part complex. See U.S. Patent No. 4,376,110. The second antibody may itself be labeled with a detectable moiety (direct sandwich assays) or may be measured using an anti-immunoglobulin antibody that is labeled with a detectable moiety (indirect sandwich assays). For example, one type of sandwich assay is an ELISA assay, in which case the detectable moiety is an enzyme.

The anti-beta secretase antibodies of the invention also are useful for *in vivo* imaging, wherein an antibody labeled with a detectable moiety is administered to an animal, preferably into the bloodstream, and the presence and location of the labeled antibody in the host is assayed. The antibody may be labeled with any moiety that is detectable in an animal, whether by nuclear magnetic resonance, radiology, or other detection means known in the art.

Antibodies of the invention may be used as therapeutics. Therapeutic antibodies are generally agonists or antagonists, in that they either enhance or reduce, respectively, at least one of the biological activities of an beta secretase polypeptide. Antagonist antibodies of the invention are antibodies or binding fragments thereof which are capable of specifically binding to an beta secretase polypeptide, fragment, variant and/or derivative, and which are capable of inhibiting or eliminating the functional activity of an beta secretase polypeptide *in vivo* or *in vitro*. In preferred embodiments, an antagonist antibody will inhibit the functional activity of an beta secretase polypeptide at least about 50 percent, and preferably at least about 80 percent. Agonist and antagonist anti-beta secretase antibodies are identified by screening assays described below.

- 53 -

Genetically Engineered Non-Human Mammals

Included within the scope of the present invention are non-human mammals such as mice, rats, rabbits, goats, or sheep in which one or both alleles of the gene encoding a native beta secretase polypeptide has (have) been disrupted ("knocked out") such that the level of expression of this gene is decreased or completely abolished. Such mammals may be prepared using techniques and methods such as those described in U.S. Patent No. 5,557,032. Such beta secretase knockout mammals have use in evaluating the effects of decreased beta secretase expression on Alzheimer's disease.

The present invention further includes non-human mammals such as mice, rats, rabbits, goats, or sheep in which the gene (or genes) encoding beta secretase polypeptides in which either the native form of the gene(s) for that mammal or a heterologous beta secretase polypeptide gene(s) is (are) over-expressed by the mammal, thereby creating a "transgenic" mammal. Such transgenic mammals may be prepared using well known methods such as those described in U.S. Patent No 5,489,743 and PCT Publication No. WO94/28122. Such transgenic mammals have use in evaluating the effects of over production of beta secretase on Alzheimer's disease.

The present invention further includes non-human mammals in which the promoter for one or more of the beta secretase polypeptides of the present invention is either activated or inactivated (using homologous recombination methods as described below) to alter the level of expression of one or more of the native beta secretase polypeptides. These mammals have uses similar to those set forth for transgenic and knockout mammals.



- 54 -

Modulators of Beta Secretase Polypeptide Activity

In some situations, it may be desirable to identify molecules that are modulators, i.e., agonists or antagonists, of the activity of beta secretase polypeptide.

Natural or synthetic molecules that modulate beta secretase can be identified using one or more of the screening assays described below. Such molecules may be administered either in an ex vivo manner, or in an in vivo manner by local or IV injection, or by oral delivery, implantation device, or the like.

The following definition is used herein for describing the assays:

"Test molecule(s)" refers to the molecule(s) that is/are under evaluation for the ability to modulate beta secretase polypeptide activity. Such test molecule may be an agonist or antagonist of beta secretase.

Methods for identifying compounds which modulate beta secretase activity are encompassed by the present invention. In general, a beta secretase polypeptide can be incubated with a test molecule under conditions which permit the interaction of the test molecule with beta secretase polypeptide, and the extent of the interaction can then be measured. The test molecule may be screened in a substantially purified form or in a crude mixture. Test molecules may be nucleic acid molecules, proteins, peptides, carbohydrates, lipids or small molecular weight organic or inorganic compounds. Once a set of test molecules has been identified as binding to an beta secretase polypeptide, the molecules may be further evaluated for their ability to increase or decrease beta secretase activity.

- 55 -

Measurement of the interaction of test molecules with beta secretase polypeptides may be carried out in several formats, including, without limitation, enzymatic assays, cell-based assays, solution-phase assays, immunoassays and *in vivo* assays. In general, test molecules are incubated with an beta secretase polypeptide for a specified period of time and the extent of beta secretase activity can then be determined by the biological activity assay set forth herein, or by other appropriate assays such as immunoassays.

The beta secretase agonist or antagonist may be a protein, peptide, carbohydrate, lipid or small molecular weight molecule which interacts with beta secretase to regulate its activity. Potential protein antagonists of beta secretase include antibodies which bind to active regions of the polypeptide and inhibit or eliminate at least once activity of beta secretase. Molecules which regulate beta secretase polypeptide expression may include nucleic acids which are complementary to nucleic acids encoding an beta secretase polypeptide, or are complementary to nucleic acids sequences which direct or control expression of beta secretase polypeptide, and which act as anti-sense regulators of expression.

In some cases, it may be desirable to evaluate two or more test compounds together for their ability to modulate beta secretase polypeptide activity. In these cases, the assays can be readily modified by adding such additional test compound(s) either simultaneous with, or subsequent to, the first test compound.

*In vitro* assays such as the biological activity assay described herein may be used advantageously to rapidly screen large numbers of compounds for effects on the activity of beta

- 56 -

secretase. The assays may be automated to screen compounds generated using phage display, synthetic peptide and chemical synthesis libraries.

Compounds which increase or decrease beta secretase activity may be screened in cell culture using cells and cell lines expressing beta secretase. Cells and cell lines may be obtained from any mammal, but preferably will be from human or other primate, canine, or rodent sources.

#### Beta Secretase Compositions and Administration

Therapeutic compositions of beta secretase polypeptides are within the scope of the present invention. Such compositions may comprise a therapeutically effective amount of a beta secretase polypeptide, fragment, variant, or derivative in admixture with a pharmaceutically acceptable agent such as a pharmaceutically acceptable carrier. The carrier material may be water for injection, preferably supplemented with other materials common in solutions for administration to mammals. Typically, a beta secretase polypeptide therapeutic compound will be administered in the form of a composition comprising purified polypeptide, fragment, variant, or derivative in conjunction with one or more physiologically acceptable agents. Neutral buffered saline or saline mixed with serum albumin are exemplary appropriate carriers. Preferably, the product is formulated as a lyophilizate using appropriate excipients (e.g., sucrose). Other standard pharmaceutically acceptable agents such as carriers, diluents, and excipients may be included as desired. Other exemplary compositions comprise Tris buffer of about pH 7.0-8.5, or acetate buffer of about pH 4.0-5.5, which may further include sorbitol or a suitable substitute therefor.

Beta secretase pharmaceutical compositions

- 57 -

typically include a therapeutically or prophylactically effective amount of beta secretase protein in admixture with one or more pharmaceutically and physiologically acceptable formulation agents selected for suitability with the mode of administration. Suitable formulation materials or pharmaceutically acceptable agents include, but are not limited to, antioxidants, preservatives, coloring, flavoring and diluting agents, emulsifying agents, suspending agents, solvents, fillers, bulking agents, buffers, delivery vehicles, diluents, excipients and/or pharmaceutical adjuvants. For example, a suitable vehicle may be water for injection, physiological saline solution, or artificial cerebrospinal fluid, possibly supplemented with other materials common in compositions for parenteral administration. Neutral buffered saline or saline mixed with serum albumin are further exemplary vehicles. The term "pharmaceutically acceptable carrier" or "physiologically acceptable carrier" as used herein refers to a formulation agent(s) suitable for accomplishing or enhancing the delivery of the beta secretase protein as a pharmaceutical composition.

The primary solvent in a composition may be either aqueous or non-aqueous in nature. In addition, the vehicle may contain other formulation materials for modifying or maintaining the pH, osmolarity, viscosity, clarity, color, sterility, stability, rate of dissolution, or odor of the formulation. Similarly, the composition may contain additional formulation materials for modifying or maintaining the rate of release of beta secretase protein, or for promoting the absorption or penetration of beta secretase protein.

The beta secretase polypeptide compositions can be administered parentally. Alternatively, the compositions may be administered intravenously or subcutaneously. When systemically administered, the

- 58 -

therapeutic compositions for use in this invention may be in the form of a pyrogen-free, parentally acceptable aqueous solution. The preparation of such pharmaceutically acceptable protein solutions, with due regard to pH, isotonicity, stability and the like, is within the skill of the art.

Therapeutic formulations of beta secretase polypeptide compositions useful for practicing the present invention may be prepared for storage by mixing the selected composition having the desired degree of purity with optional physiologically acceptable carriers, excipients, or stabilizers (*Remington's Pharmaceutical Sciences*, 18th Edition, A.R. Gennaro, ed., Mack Publishing Company [1990]) in the form of a lyophilized cake or an aqueous solution. Acceptable carriers, excipients or stabilizers are nontoxic to recipients and are preferably inert at the dosages and concentrations employed, and include buffers such as phosphate, citrate, or other organic acids; antioxidants such as ascorbic acid; low molecular weight polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as Tween, pluronics or polyethylene glycol (PEG).

The optimal pharmaceutical formulation will be determined by one skilled in the art depending upon the intended route of administration, delivery format and desired dosage. See for example, *Remington's Pharmaceutical Sciences*, 18th Ed. (1990, Mack Publishing Co., Easton, PA 18042 pages 1435-1712).

- 59 -

Such compositions may influence the physical state, stability, rate of *in vivo* release, and rate of *in vivo* clearance of the present beta secretase protein.

An effective amount of an beta secretase polypeptide composition to be employed therapeutically will depend, for example, upon the therapeutic objectives such as the indication for which the beta secretase polypeptide is being used, the route of administration, and the condition of the patient. Accordingly, it may be necessary for the therapist to titer the dosage and modify the route of administration as required to obtain the optimal therapeutic effect. A typical daily dosage may range from about 0.1 µg/kg to up to 100 mg/kg or more, depending on the factors mentioned above. Typically, a clinician will administer the composition until a dosage is reached that achieves the desired effect. The composition may therefore be administered as a single dose, or as two or more doses (which may or may not contain the same amount of beta secretase polypeptide) over time, or as a continuous infusion via implantation device or catheter.

As further studies are conducted, information will emerge regarding appropriate dosage levels for treatment of various conditions in various patients, and the ordinary skilled worker, considering the therapeutic context, the type of disorder under treatment, the age and general health of the recipient, will be able to ascertain proper dosing.

The beta secretase polypeptide composition to be used for *in vivo* parenteral administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes. Where the composition is lyophilized, sterilization using these

5

- 60 -

10

methods may be conducted either prior to, or following, lyophilization and reconstitution. The composition for parenteral administration ordinarily will be stored in lyophilized form or in solution.

15

5       Therapeutic compositions generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

20

Effective administration forms, such as (1) 10 slow-release formulations, (2) inhalant mists, or (3) orally active formulations are also envisioned. The beta secretase pharmaceutical composition also may be formulated for parenteral administration. Such parenterally administered therapeutic compositions are 15 typically in the form of a pyrogen-free, parenterally acceptable aqueous solution comprising beta secretase in a pharmaceutically acceptable vehicle. The beta secretase pharmaceutical compositions also may include 20 particulate preparations of polymeric compounds such as polylactic acid, polyglycolic acid, etc. or the introduction of beta secretase into liposomes. Hyaluronic acid may also be used, and this may have the 25 effect of promoting sustained duration in the circulation.

30

35

40

25       A particularly suitable vehicle for parenteral injection is sterile distilled water in which beta secretase is formulated as a sterile, isotonic solution, properly preserved. Yet another preparation may involve the formulation of beta 30 secretase with an agent, such as injectable microspheres, bio-erodible particles or beads, or liposomes, that provides for the controlled or 45 sustained release of the protein product which may then be delivered as a depot injection. Other suitable 35 means for the introduction of beta secretase include 50 implantable drug delivery devices which contain the

55

55

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

B14

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification:</b> C12N 15/52, A01K 67/027, A61K 38/48, A61K 48/00, C07K 16/40, C12N 9/64, C12N 15/62, G01N 33/566	<b>A1</b>	<b>(11) International Publication Number:</b> WO 00/58479 <b>(43) International Publication Date:</b> 05 October 2000 (05.10.2000)
<b>(21) International Application Number:</b> PCT/US00/07755 <b>(22) International Filing Date:</b> 23 March 2000 (23.03.2000) <b>(30) Priority Data:</b> 09/277,229 26 March 1999 (26.03.1999) US <b>(60) Parent Application or Grant</b> AMGEN INC. [/]; (). CITRON, Martin [/]; (). VASSAR, Robert, James [/]; (). BENNETT, Brian, Drake [/]; (). ODRE, Steven, M. ; ().		<b>Published</b>
<b>(54) Title: BETA SECRETASE GENES AND POLYPEPTIDES</b> <b>(54) Titre: GENES ET POLYPEPTIDES BETA SECRETASES</b>		
<b>(57) Abstract</b> Disclosed are novel genes encoding beta secretase polypeptides. Also disclosed are methods of making and using the polypeptides. <b>(57) Abrégé</b> L'invention concerne de nouveaux gènes codant pour des polypeptides bêta sécrétase, ainsi que des méthodes de fabrication et d'utilisation des polypeptides.		



- 61 -

beta secretase.

The preparations of the present invention may include other components, for example parenterally acceptable preservatives, tonicity agents, cosolvents,

wetting agents, complexing agents, buffering agents, antimicrobials, antioxidants and surfactants, as are well known in the art. For example, suitable tonicity enhancing agents include alkali metal halides

(preferably sodium or potassium chloride), mannitol, sorbitol and the like. Suitable preservatives include, but are not limited to, benzalkonium chloride, thimerosal, phenethyl alcohol, methylparaben, propylparaben, chlorhexidine, sorbic acid and the like. Hydrogen peroxide may also be used as preservative.

Suitable cosolvents are for example glycerin, propylene glycol and polyethylene glycol. Suitable complexing agents are for example caffeine, polyvinylpyrrolidone, beta-cyclodextrin or hydroxypropyl-beta-cyclodextrin.

Suitable surfactants or wetting agents include sorbitan esters, polysorbates such as polysorbate 80, tromethamine, lecithin, cholesterol, tyloxapal and the like. The buffers can be conventional buffers such as borate, citrate, phosphate, bicarbonate, or Tris-HCl.

The formulation components are present in concentration that are acceptable to the site of administration. For example, buffers are used to maintain the composition at physiological pH or at slightly lower pH, typically within a pH range of from about 5 to about 8.

A pharmaceutical composition may be formulated for inhalation. For example, beta secretase may be formulated as a dry powder for inhalation. beta secretase inhalation solutions may also be formulated in a liquefied propellant for aerosol delivery. In yet another formulation, solutions may be nebulized.

It is also contemplated that certain

- 62 -

formulations containing beta secretase may be administered orally. Beta secretase which is administered in this fashion may be formulated with or without those carriers customarily used in the compounding of solid dosage forms such as tablets and capsules. For example, a capsule may be designed to release the active portion of the formulation at the point in the gastrointestinal tract when bioavailability is maximized and pre-systemic degradation is minimized. Additional agents may be included to facilitate absorption of beta secretase. Diluents, flavorings, low melting point waxes, vegetable oils, lubricants, suspending agents, tablet disintegrating agents, and binders may also be employed.

Another preparation may involve an effective quantity of beta secretase in a mixture with non-toxic excipients which are suitable for the manufacture of tablets. By dissolving the tablets in sterile water, or other appropriate vehicle, solutions can be prepared in unit dose form. Suitable excipients include, but are not limited to, inert diluents, such as calcium carbonate, sodium carbonate or bicarbonate, lactose, or calcium phosphate; or binding agents, such as starch, gelatin, or acacia; or lubricating agents such as magnesium stearate, stearic acid, or talc.

Additional beta secretase formulations will be evident to those skilled in the art, including formulations involving beta secretase in combination with one or more other therapeutic agents. Techniques for formulating a variety of other sustained- or controlled-delivery means, such as liposome carriers, bio-erodible microparticles or porous beads and depot injections, are also known to those skilled in the art. See, for example, the Supersaxo et al. description of controlled release porous polymeric microparticles for

- 63 -

the delivery of pharmaceutical compositions  
(International Publication No. WO 93/15722;  
International Application No. PCT/US93/00829) the  
disclosure of which is hereby incorporated by  
reference.

Regardless of the manner of administration,  
the specific dose may be calculated according to body  
weight, body surface area or organ size. Further  
refinement of the calculations necessary to determine  
the appropriate dosage for treatment involving each of  
the above mentioned formulations is routinely made by  
those of ordinary skill in the art and is within the  
ambit of tasks routinely performed by them.  
Appropriate dosages may be ascertained through use of  
appropriate dose-response data.

The route of administration of the  
composition is in accord with known methods, e.g. oral,  
injection or infusion by intravenous, intraperitoneal,  
intracerebral (intraparenchymal),  
intracerebroventricular, intramuscular, intraocular,  
intraarterial, or intralesional routes, or by sustained  
release systems or implantation device which may  
optionally involve the use of a catheter. Where  
desired, the compositions may be administered  
continuously by infusion, bolus injection or by  
implantation device.

Alternatively or additionally, the  
composition may be administered locally via  
implantation into the affected area of a membrane,  
sponge, or other appropriate material on to which beta  
secretase polypeptide has been absorbed or  
encapsulated.

Where an implantation device is used, the  
device may be implanted into any suitable tissue or  
organ, and delivery of beta secretase polypeptide may  
be directly through the device via bolus, or via

- 64 -

continuous administration, or via catheter using continuous infusion.

Beta secretase polypeptide may be administered in a sustained release formulation or preparation. Suitable examples of sustained-release preparations include semipermeable polymer matrices in the form of shaped articles, e.g. films, or microcapsules. Sustained release matrices include polyesters, hydrogels, polylactides (U.S. 3,773,919, EP 58,481), copolymers of L-glutamic acid and gamma ethyl-L-glutamate (Sidman et al, *Biopolymers*, 22: 547-556 [1983]), poly (2-hydroxyethyl-methacrylate) (Langer et al., *J. Biomed. Mater. Res.*, 15: 167-277 [1981] and Langer, *Chem. Tech.*, 12: 93-105 [1982]), ethylene vinyl acetate (Langer et al., *supra*) or poly-D(-)-3-hydroxybutyric acid (EP 133,989). Sustained-release compositions also may include liposomes, which can be prepared by any of several methods known in the art (e.g., Eppstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688-3692 [1985]; EP 36,676; EP 88,045; EP 143,949).

The beta secretase polypeptides, fragments thereof, variants, and derivatives, may be employed alone, together, or in combination with other pharmaceutical compositions. The beta secretase polypeptides, fragments, variants, and derivatives may be used in combination with other medicinal compounds such as, for example, cytokines, growth factors, antibiotics, anti-inflammatories, and/or chemotherapeutic agents as is appropriate for the indication being treated.

In some cases, it may be desirable to use beta secretase polypeptide compositions in an ex vivo manner. Here, cells, tissues, or organs that have been removed from the patient are exposed to beta secretase polypeptide compositions after which the cells, tissues

- 65 -

and/or organs are subsequently implanted back into the patient.

In other cases, an beta secretase polypeptide may be delivered through implanting into patients certain cells that have been genetically engineered, using methods such as those described herein, to express and secrete the polypeptides, fragments, variants, or derivatives. Such cells may be animal or human cells, and may be derived from the patient's own tissue or from another source, either human or non-human. Optionally, the cells may be immortalized. However, in order to decrease the chance of an immunological response, it is preferred that the cells be encapsulated to avoid infiltration of surrounding tissues. The encapsulation materials are typically biocompatible, semi-permeable polymeric enclosures or membranes that allow release of the protein product(s) but prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissues.

Methods used for membrane encapsulation of cells are familiar to the skilled artisan, and preparation of encapsulated cells and their implantation in patients may be accomplished without undue experimentation. See, e.g., U.S. Patent Nos. 4,892,538; 5,011,472; and 5,106,627. A system for encapsulating living cells is described in PCT WO 91/10425 (Aebischer et al.). The cells, with or without encapsulation, may be implanted into suitable body tissues or organs of the patient.

#### Gene Therapy and Homologous Recombination

Further included in the scope of the present invention is production of beta secretase polypeptide by homologous recombination, and production of beta secretase polypeptide using control elements introduced

- 66 -

into cells containing beta secretase DNA. For example, homologous recombination methods may be used to modify a cell containing a transcriptionally active beta secretase gene to produce a cell which does not express therapeutically efficacious amounts of beta secretase.

Homologous recombination is a technique originally developed for targeting genes to induce or correct mutations in transcriptionally active genes (see Kucherlapati, *Prog. in Nucl. Acid Res. and Mol. Biol.*, 36:301, 1989). The basic technique was developed as a method for introducing specific mutations into specific regions of the mammalian genome (Thomas et al., *Cell*, 44:419-428, 1986; Thomas et al., *Cell*, 51:503-512, 1987; Doetschman et al., *Proc. Natl. Acad. Sci.*, 85:8583-8587, 1988) or to correct specific mutations within defective genes (Doetschman et al., *Nature*, 330:576-578, 1987). Exemplary homologous recombination techniques are described in U.S. 5,272,071 (EP 91 90 3051, EP Publication No. 505 500; and International Publication No. WO 91/09955).

Through homologous recombination, the DNA sequence to be inserted into the genome can be directed to a specific region of the gene of interest by attaching it to targeting DNA. The targeting DNA is a nucleic acid molecule that is complementary to a region of the genomic DNA. Small pieces of targeting DNA that are complementary to a specific region of the genome are put in contact with the parental strand during the DNA replication process. It is a general property of DNA that has been inserted into a cell to hybridize, and therefore, recombine with other pieces of endogenous DNA through shared homologous regions. If this complementary strand is attached to an oligonucleotide that contains a mutation or a different sequence or an additional nucleotide, it too is incorporated into the newly synthesized strand as a

- 67 -

result of the recombination. As a result of the proofreading function, it is possible for the new DNA to serve as the template. Thus, this new DNA is incorporated into the genome.

Attached to these pieces of targeting DNA are regions of DNA which may interact with the expression of a beta secretase protein. For example, a promoter/enhancer element, a suppresser, or an exogenous transcription modulatory element is inserted in the genome of the intended host cell in proximity and orientation sufficient to influence the transcription of DNA encoding the desired beta secretase protein. The control element controls a portion of the DNA present in the host cell genome. Thus, the expression of beta secretase protein may be achieved not by transfection of DNA that encodes the beta secretase gene itself, but rather by the use of targeting DNA (containing regions of homology with the endogenous gene of interest) coupled with DNA regulatory segments that provide the endogenous gene sequence with recognizable signals for transcription of a beta secretase protein.

In an exemplary method, expression of a desired targeted gene in a cell (i.e., a desired endogenous cellular gene) can be altered by introducing an exogenous DNA molecule into a preselected site of genomic DNA. The exogenous DNA molecule typically includes at least a regulatory sequence, an exon and a splice donor site. This exogenous DNA is introduced into the genomic DNA in a location so as to produce a new transcription unit in which the regulatory sequence, the exon and the splice donor site present in the exogenous DNA are operatively linked to the endogenous gene. As a result of introduction of the exogenous DNA into the genomic DNA, the expression of the desired endogenous gene is altered.

- 68 -

Altered gene expression, as used herein, encompasses activating (or causing to be expressed) a gene which is normally silent (unexpressed) in the cell as obtained, increasing expression of a gene which is not expressed at physiologically significant levels in the cell as obtained, changing the pattern of regulation or induction such that it is different than occurs in the cell as obtained, and reducing (including eliminating) expression of a gene which is expressed in the cell as obtained.

The present invention further relates to DNA constructs useful in the method of altering expression of a target gene. Exemplary DNA constructs typically comprise the following components: (a) a targeting sequence; (b) a regulatory sequence; (c) an exon; and (d) an unpaired splice-donor site. The targeting sequence in the DNA construct directs the integration of elements (a) - (d) into a target gene in a cell such that the elements (b) - (d) are operatively linked to sequences of the endogenous target gene. In another embodiment, the DNA constructs comprise: (a) a targeting sequence, (b) a regulatory sequence, (c) an exon, (d) a splice-donor site, (e) an intron, and (f) a splice-acceptor site, wherein the targeting sequence directs the integration of elements (a) - (f) such that the elements of (b) - (f) are operatively linked to the endogenous gene. The targeting sequence is homologous to the preselected site in the cellular chromosomal DNA with which homologous recombination is to occur. In the construct, the exon is generally 3' of the regulatory sequence and the splice-donor site is 3' of the exon.

If the sequence of a particular gene is known, such as the nucleic acid sequence of beta secretase presented herein, a piece of DNA that is complementary to a selected region of the gene can be



- 69 -

synthesized or otherwise obtained, such as by  
appropriate restriction of the native DNA at specific  
recognition sites bounding the region of interest.  
This piece serves as a targeting sequence upon  
insertion into the cell and will hybridize to its  
homologous region within the genome. If this  
hybridization occurs during DNA replication, this piece  
of DNA, and any additional sequence attached thereto,  
will act as an Okazaki fragment and will be  
backstitched into the newly synthesized daughter strand  
of DNA. The present invention, therefore, includes  
nucleotides encoding a beta secretase molecule, which  
nucleotides may be used as targeting sequences.

Cell based therapy via genetic manipulation  
is also included in the scope of the present invention.  
In those situations where it is desirable to increase  
the level of beta secretase activity in cells, beta  
secretase cell therapy, e.g., implantation of cells  
producing beta secretase, may be appropriate. This  
embodiment encompasses implanting into patients cells  
capable of synthesizing and secreting a biologically  
active form of beta secretase. Such beta secretase-  
producing cells may be cells that are natural producers  
of beta secretase or may be recombinant cells whose  
ability to produce beta secretase has been augmented by  
transformation with a gene encoding the desired beta  
secretase molecule or with a gene augmenting the  
expression of beta secretase. Such a modification may  
be accomplished by means of a vector suitable for  
delivering the gene as well as promoting its expression  
and secretion. In order to minimize a potential  
immunological reaction in patients being administered a  
beta secretase protein or polypeptide of a foreign  
species, it is preferred that the natural cells  
producing beta secretase be of human origin and produce  
human beta secretase. Likewise, it is preferred that

- 70 -

the recombinant cells producing beta secretase be transformed with an expression vector containing a gene encoding a human beta secretase molecule.

Implanted cells may be encapsulated to avoid infiltration of surrounding tissue. Human or non-human animal cells may be implanted in patients in biocompatible, semipermeable polymeric enclosures or membranes that allow release of beta secretase, but that prevent destruction of the cells by the patient's immune system or by other detrimental factors from the surrounding tissue. Alternatively, the patient's own cells, transformed to produce beta secretase *ex vivo*, could be implanted directly into the patient without such encapsulation.

Techniques for the encapsulation of living cells are known in the art, and the preparation of the encapsulated cells and their implantation in patients may be accomplished without undue experimentation. For example, Baetge *et al.* (International Publication No. WO 95/05452) describe membrane capsules containing genetically engineered cells for the effective delivery of biologically active molecules. The capsules are biocompatible and are easily retrievable. The capsules encapsulate cells transfected with recombinant DNA molecules comprising DNA sequences coding for biologically active molecules operatively linked to promoters that are not subject to down regulation *in vivo* upon implantation into a mammalian host. The devices provide for the delivery of the molecules from living cells to specific sites within a recipient. In addition, see U.S. Patent Numbers 4,392,538, 5,011,472, and 5,106,527. A system for encapsulating living cells is described in PCT Application WO 91/10425 of Aebischer *et al.* See also PCT Application WO 91/10470 of Aebischer *et al.*; Winn *et al.*, *Exper. Neurol.*, 113:322-329, 1991; and Aebischer *et al.*, *Exper.*

- 71 -

Neurol., 111:269-275, 1991.

*In vivo* and *in vitro* gene therapy delivery of beta secretase is also envisioned. *In vivo* gene therapy may be accomplished by introducing the gene encoding beta secretase into cells via local injection of a polynucleotide molecule or other appropriate delivery vectors. (Hefti, *J. Neurobiology*, 25:1418-1435, 1994). For example, a polynucleotide molecule encoding beta secretase may be contained in an adeno-associated virus vector for delivery to the targeted cells (e.g., Johnson, International Publication No. WO 95/34670; International Application No. PCT/US95/07178). The recombinant adeno-associated virus (AAV) genome typically contains AAV inverted terminal repeats flanking a DNA sequence encoding beta secretase operably linked to functional promoter and polyadenylation sequences.

Alternative viral vectors include, but are not limited to, retrovirus, adenovirus, herpes simplex virus and papilloma virus vectors. U.S. 5,672,344 (issued September 30, 1997, Kelley et al., University of Michigan) describes an *in vivo* viral-mediated gene transfer system involving a recombinant neurotrophic HSV-1 vector. U.S. 5,399,346 (issued March 21, 1995, Anderson et al., Department of Health and human Services) provides examples of a process for providing a patient with a therapeutic protein by the delivery of human cells which have been treated *in vitro* to insert a DNA segment encoding a therapeutic protein. Additional methods and materials for the practice of gene therapy techniques are described in U.S. 5,631,236 (issued May 20, 1997, Woo et al., Baylor College of Medicine) involving adenoviral vectors; U.S. 5,672,510 (issued September 30, 1997, Eglitis et al., Genetic Therapy, Inc.) involving retroviral vectors; and U.S. 5,635,399 (issued June 3, 1997, Kriegler et al., Chiron

- 72 -

Corporation) involving retroviral vectors expressing cytokines.

Nonviral delivery methods include liposome-mediated transfer, naked DNA delivery (direct injection), receptor-mediated transfer (ligand-DNA complex), electroporation, calcium phosphate precipitation and microparticle bombardment (e.g., gene gun). Gene therapy materials and methods may also include inducible promoters, tissue-specific enhancer-promoters, DNA sequences designed for site-specific integration, DNA sequences capable of providing a selective advantage over the parent cell, labels to identify transformed cells, negative selection systems and expression control systems (safety measures), cell-specific binding agents (for cell targeting), cell-specific internalization factors, transcription factors to enhance expression by a vector as well as methods of vector manufacture. Such additional methods and materials for the practice of gene therapy techniques are described in U.S. 4,970,154 (issued November 13, 1990, D.C. Chang, Baylor College of Medicine) electroporation techniques; WO 9640953 (published 961219, Smith et al., Baylor College of Medicine) nuclear ligands; U.S. 5,679,559 (issued October 21, 1997, Kim et al., University of Utah Research Foundation) concerning a lipoprotein-containing system for gene delivery; U.S. 5,676,954 (issued October 14, 1997, K.L. Brigham, Vanderbilt University involving liposome carriers; U.S. 5,593,875 (issued January 14, 1997, Wurm et al., Genentech, Inc.) concerning methods for calcium phosphate transfection; and U.S. 4,945,050 (issued July 31, 1990, Sanford et al., Cornell Research Foundation) wherein biologically active particles are propelled at cells at a speed whereby the particles penetrate the surface of the cells and become incorporated into the interior of the cells.

- 73 -

Expression control techniques include chemical induced regulation (e.g., WO 9641865 and WO 9731899), the use of a progesterone antagonist in a modified steroid hormone receptor system (e.g., U.S. 5,364,791), ecdysone control systems (e.g., WO 9637609), and positive tetracycline-controllable transactivators (e.g., U.S. 5,589,362; U.S. 5,650,298; and U.S. 5,654,168).

It is also contemplated that beta secretase gene therapy or cell therapy can further include the delivery of a second polypeptide. For example, the host cell may be modified to express and release both beta secretase and a second polypeptide of interest. Alternatively, the beta secretase and the second polypeptide of interest may be expressed in and released from separate cells. Such cells may be separately introduced into the patient or the cells may be contained in a single implantable device, such as the encapsulating membrane described above.

Gene therapy can be used to decrease beta secretase polypeptide expression by modifying the nucleotide sequence of the endogenous promoter(s). Such modification is typically accomplished via homologous recombination methods. For example, a DNA molecule containing all or a portion of the promoter of the beta secretase gene(s) selected for inactivation can be engineered to remove and/or replace pieces of the promoter that regulate transcription. Here, the TATA box and/or the binding site of a transcriptional activator of the promoter may be deleted using standard molecular biology techniques; such deletion can inhibit promoter activity thereby repressing transcription of the corresponding beta secretase gene. Deletion of the TATA box or transcription activator binding site in the promoter may be accomplished by generating a DNA construct comprising all or the relevant portion of the

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

beta secretase polypeptide promoter(s) (from the same or a related species as the beta secretase gene(s) to be regulated) in which one or more of the TATA box and/or transcriptional activator binding site nucleotides are mutated via substitution, deletion and/or insertion of one or more nucleotides such that the TATA box and/or activator binding site has decreased activity or is rendered completely inactive. This construct, which also will typically contain at least about 500 bases of DNA that correspond to the native (endogenous) 5' and 3' DNA sequences adjacent to the promoter segment that has been modified, may be introduced into the appropriate cells (either ex vivo or in vivo) either directly or via a viral vector as described above. Typically, integration of the construct into the genomic DNA of the cells will be via homologous recombination, where the 5' and 3' DNA sequences in the promoter construct can serve to help integrate the modified promoter region via hybridization to the endogenous chromosomal DNA.

Other gene therapy methods may also be employed where it is desirable to inhibit the activity of one or more beta secretase polypeptides. For example, antisense DNA or RNA molecules, which have a sequence that is complementary to at least a portion of the selected beta secretase polypeptide gene(s) can be introduced into the cell. Typically, each such antisense molecule will be complementary to the start site (5' end) of each selected beta secretase gene. When the antisense molecule then hybridizes to the corresponding beta secretase mRNA, translation of this mRNA is prevented.

Alternatively, gene therapy may be employed to create a dominant-negative inhibitor of one or more beta secretase polypeptides. In this situation, the DNA encoding a mutant full length or truncated

- 75 -

polypeptide of each selected beta secretase polypeptide can be prepared and introduced into the cells of a patient using either viral or non-viral methods as described above. Each such mutant is typically designed to compete with endogenous polypeptide in its biological role.

#### Utility

Nucleic acid molecules of the invention may be used to map the locations of the beta secretase gene and related genes on chromosomes. Mapping may be done by techniques known in the art, such as PCR amplification and *in situ* hybridization.

In addition, the nucleic acid molecules can be used as anti-sense inhibitors of beta secretase expression. Such inhibition may be effected by nucleic acid molecules which are complementary to and hybridize to expression control sequences (triple helix formation) or to beta secretase mRNA. Anti-sense probes may be designed by available techniques using the sequence of beta secretase disclosed herein. Anti-sense inhibitors provide information relating to the decrease or absence of an beta secretase polypeptide in a cell or organism.

Hybridization probes may be prepared using the beta secretase nucleic acid molecules provided herein to screen cDNA, genomic or synthetic DNA libraries for related sequences. Regions of the DNA and/or amino acid sequence of beta secretase that exhibit significant identity to known sequences are readily determined using sequence alignment algorithms disclosed above and those regions may be used to design probes for screening.

Further, the nucleic acid molecules of the invention may be used for gene therapy. Nucleic acid molecules which express beta secretase *in vivo* provide

- 76 -

information relating to the effects of the polypeptide in cells or organisms.

Beta secretase nucleic acid molecules, fragments, and/or derivatives that do not themselves encode biologically active polypeptides may be useful as hybridization probes in diagnostic assays to test, either qualitatively or quantitatively, for the presence of beta secretase DNA or corresponding RNA in mammalian tissue or bodily fluid samples.

Beta secretase polypeptide fragments, variants, and/or derivatives, whether biologically active or not, are useful for preparing antibodies that bind to a beta secretase polypeptide. The antibodies may be used for *in vivo* and *in vitro* diagnostic purposes, such as in labeled form to detect the presence of beta secretase polypeptide in a body fluid or cell sample. Diagnosis of conditions such as Alzheimer's disease, Down's syndrome, and amyloid angiopathy may be accomplished by measuring beta-secretase levels and activity from bodily tissue or cell samples such as plasma, and comparing this data against known "normal" standards for each such condition. Beta-secretase levels can be measured by ELISA or Western blot for example; beta-secretase activity can be measured by the assays set forth in the Examples below. Abnormal results of one or both assays could indicate that a person is at high risk of developing such diseases, and early treatment could be started via administration of a beta-secretase inhibitor, for example.

Based on Northern blot data presented in the Examples below, beta-secretase expression is relatively high in the pancreas. This suggests that beta-secretase may be implicated in diseases such as acute and/or chronic pancreatitis, pancreatic cancer, and pancreobiliary duct obstruction, and modulation of the



- 77 -

activity of beta-secretase may decrease the extent of, or completely prevent, such disorders.

The beta-secretase antibodies may bind to a beta-secretase polypeptide so as to diminish or block at least one activity characteristic of beta secretase polypeptide, or may bind to a beta-secretase polypeptide to increase its activity.

A plasmid containing cDNA encoding full length beta secretase has been transformed into in *E. coli* cells, and the cells have been deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209 on March 11, 1999. The cells have been designated as accession no. 207159.

A plasmid containing cDNA encoding amino acids 1-460 of beta-secretase fused to the Fc portion of human IgG beta has been transformed into in *E. coli* cells, and the cells have been deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, VA 20110-2209 on March 11, 1999. The cells have been designated as accession no. 207158.

The following examples are intended for illustration purposes only, and should not be construed as limiting the scope of the invention in any way.

- 73 -

Example I: Production of an APP Expressing Cell Line

The human APP-695 cDNA containing the Swedish mutation (referred to as "APPANL" the sequence of which has been published Mullan et al., *Nature Genetics* 1: 345-347 (1992)) was digested with the restriction enzymes *HindIII* and *NotI*, and the approximately 2.6 kb *HindIII* - *Not* fragment was inserted into the vector pCMVi (Cell & Molecular Technologies Inc; Lavallette, NJ) to generate the vector pCMVi-APPANL).

Human embryonic kidney 293 cells containing the SV40 large T antigen (Cell & Molecular Technologies, Lavallette, NJ) were maintained in high glucose Dulbecco's modified Eagle's medium supplemented with about 10 percent fetal calf serum. At approximately 70 percent confluence the cells were co-transfected with pCMVi-APPANL and pRSV-Puro (Cell & Molecular Technologies Inc; Lavallette, NJ) using the standard calcium phosphate method (Gorman et al, *DNA Prot. Eng. Tech.* 2: 3-10 (1990)). After about forty eight hours, puromycin resistant colonies were selected by adding puromycin (Sigma Inc., St. Louis, MO) at about 5 micrograms/ml to the culture medium. Fourteen days later, individual resistant colonies were isolated, using a pipettman P200 (Rainin, Emeryville, CA) expanded into a 24 well dish and allowed to grow

- 79 -

confluent. The conditioned media were analyzed for the presence of secreted APP by Western dot blot using the monoclonal antibody 22C11 (Boehringer Mannheim Corp., Indianapolis, IN).

One colony was grown up and a cell line referred to as "293T-APPsw101" was generated from it.

Example II: Construction of a Human Embryonic Kidney Cell Expression Library

Approximately 10<sup>6</sup> 293 human embryonic kidney cells (ATCC CRL-1573), which are known to express the beta secretase gene, were grown to about 80 percent confluency in DMEM high glucose media supplemented with about 10 percent fetal bovine serum using T225 tissue culture flasks (Corning/Costar, Encinitas, CA). Cells were harvested in about 3 ml of 0.5 mM EDTA in PBS and pelleted at about 1100 RPM in a Beckman table top centrifuge. The cells were washed twice with cold PBS and snap frozen in liquid nitrogen as a dry cell pellet. Poly A+ mRNA was harvested from the cell pellet using the FastTrack™ method of mRNA isolation (Invitrogen, San Diego CA) according to the manufacturer's protocol. An oligo dT primed, directional cDNA expression library was prepared from about three µg of this poly A+ mRNA using the Superscript™ plasmid system for cDNA synthesis (Gibco

- 80 -

BRL, Gaithersburg, MD) according to the manufacturer's protocol. *Sal I* adaptors were ligated to the double stranded cDNA following the manufacturer's instructions.

After adaptor ligation, the cDNA was digested to completion with *Not I*, and size fractionated as per the manufacturer's protocol by column chromatography over a Sephacryl™ S-500 column which excludes cDNAs of 500 bp and smaller (provided by the manufacturer). Column fractions were counted on a Beckman scintillation counter and Cherenkov counts were obtained for each fraction. cDNAs from fractions 10-12 were ethanol precipitated and resuspended in about 22 microliters of nuclease free water. The cDNA products which averaged about 2.0 kb in length were directionally ligated into a CMV based expression vector (comparable to pRK-5; Pharmingen Inc., San Diego, CA) which had been previously digested with *Not I* and *Sal I*. The ligated cDNA was then introduced into electrocompetent ElectroMax™ DH1CB *E. coli* cells (Gibco BRL, Gaithersburg, MD) using standard electroporation procedures. The cDNA library was titered by serial dilution of the transformation cell mixture.

The cDNA library was subdivided into pools of 100, plated on LB agar plates containing about 100 micrograms/ml Ampicillin, and allowed to grow for about twenty hours. Cell colonies from each independent

- 81 -

5  
10  
15  
20  
25  
30  
35  
40  
45  
50  
55

plate were then harvested in about two ml of 2xYT media, transferred to deep-well 96 well dishes and allowed to grow for an additional five to six hours at 37° C. The cells were pelleted by centrifugation at 3000 RPM in a Beckman table top centrifuge and DNA was prepared using the TurboPrep™ method of DNA isolation (Qiagen Inc., Chatsworth, CA) according to the manufacturer's protocol. DNA from each pool was arrayed in 96 well Corning Costar UV plates (CORNING Costar, Charlotte, NC) and quantitated by O.D. 260 using a Molecular Devices Spectra Max plus (Molecular Dynamics, Sunnyvale, CA).

A duplicate of each master cDNA 96 well plate was prepared. Each well of the duplicate plate contained about 75 microliters of Optimem I medium (Life Technologies Inc, Gaithersburg, MD) and about 300 ng of cDNA library. About 75 microliters of a mixture of 12 microliters DMRIE-C Lipid per ml of Optimem I medium (Life Technologies Inc., Gaithersburg, MD) was added to each well of the 96 well plate using a Multipette (Sagian Inc. Indianapolis, IN) robotic pipettor. The DMRIE C was allowed to complex for about 30 minutes. Meanwhile, the media was removed from a 96 well dish in which each well had been plated with about 15,000 cells/well of the 293T-APPsw 101 cell line two days before. About 125 microliters of the DNA/Lipid complex was added to each well of cells. The cells

- 82 -

containing the DNA/lipid mixture were then incubated at 37 degrees C, 5 percent CO<sub>2</sub> for about five hours. After incubation, 125 µl Optimem I supplemented with 20 percent FBS and 2x Penicillin/Streptomycin (0.1 unit/ml Penicillin, 0.1 µg/ml Streptomycin) (Life Technologies Inc., Gaithersburg, MD) was added to stop the transfection. The following morning, the media was removed and a fresh medium containing high glucose DMEM, 10 percent FBS, and Pen/Strep was added, and the cells were allowed to recover for about 24 hours. Approximately 48 hours post transfection, fresh media was added and the cells were incubated for 6-8 hours. The medium from each well was then harvested, diluted 1:1 in Superblock (Pierce Inc., Rockford, IL), stamped into pre-coated ELISA plates and analyzed for the presence of A-beta as described below.

Example III: Beta-Secretase Activity Assays With Cultured Cells

Beta secretase generates two forms of the A-beta peptide, a 40 amino acid form and a 42 amino acid form. In most people, about ninety percent of the A-beta is the 40 amino acid form, and about 10 percent is the 42 amino acid form. However, beta-amyloid plaques consist primarily of the 42 amino acid form.

- 83 -

and in those individuals with familial Alzheimer's disease, there is a larger proportion of the 42 amino acid form as compared with the 40 amino acid form.

To analyze the level of A-beta in the cell culture medium obtained from each well of each plate, the following two assays were conducted on each medium.

A. Assay for Total Beta Amyloid Peptide

A monoclonal antibody against amino acids

17-24 of A-beta (Senetek Inc., St Louis, MO, item number 220-10) referred to as m4G8, was diluted to a concentration of about 150 micrograms/ml in coating buffer (1.59g/l Na<sub>2</sub>CO<sub>3</sub>, 2.93g/l NaHCO<sub>3</sub>, pH 9.69). About ninety microliters per well of coating buffer was then dispensed on to a polystyrene high binding 96 well flat-bottomed plate (Corning Costar, New York, NY; item number 3590). About ten microliters per well of the m4G8 antibody diluted to about 150 micrograms/ml were added to the plate coating resulting in a final concentration of about 1.5 micrograms per well of m4G8. Each plate is then incubated overnight at 4C on a plate mixer. The plates were then washed with wash buffer (7.1108g/l Tris-HCl: C<sub>4</sub>H<sub>9</sub>NO<sub>2</sub>-HCl, 0.5984g/l Tris Base: C<sub>4</sub>H<sub>9</sub>NO<sub>2</sub>, 8.766g NaCl, 0.5ml/l polyoxyethylene sorbitan monolaurate: C<sub>12</sub>H<sub>25</sub>CO<sub>2</sub>) four times, and the non-specific binding sites were blocked with 300 microliters per well of Superblock TBS (Pierce Chemical Company,

- 84 -

Rockford, IL 61105 item number 3753522) for about three hours at room temperature.

The samples used for calibration were prepared from a commercially obtained solution of Ab<sub>1-42</sub>, (5 micrograms/ml in DMSO; Quality Control Biochemicals, Inc. MA). The most concentrated sample was a 40,000 pg/ml solution, and was prepared by dilution into Superblock TBS (Pierce Chemical CO., Rockford, IL 61105). Sequential dilutions were then made in diluent to obtain 20,000, 10,000, 5,000, 2,500, 1,250, 625, 312.5, 156.25, 78.125 and 39.06 pg/ml.

One hundred microliters per well calibrators or 2.5 microliters per well of samples were applied to each well of each the microtiter plate; to each well was previously added one hundred microliters Superblock TBS plus m4G8 antibody. The plates were covered and incubated at about 4C overnight with mixing after which they were washed four times with wash buffer at about 4C.

To assess the amount of A-beta in each sample, the antibody m6E10-biotin raised against residues 1-17 of A-beta (Senetek Inc., St Louis, MO, item number 340-10), was diluted to a final concentration of about 0.5 micrograms/ml into antibody diluent (Superblock TBS as described above) containing 2 percent normal mouse serum (Sigma Chemical Co., St. Louis, MO; item number S-3509) and 2 percent normal



- 85 -

5  
10 goat serum (Gibco/BRL, Grand Island, NY; item number 15210-064). About 100 microliters of this diluted antibody were added to each well. The plate was covered and incubated for about 2 hours at 4C with  
15 5 mixing. The plate was then washed four times with wash buffer at 4C.

20 Europium labeled streptavidin (Wallac; Gaithersburg, MD; item number 1244-360) was diluted 1:1000 in Assay Buffer (Wallac; item number 1244-106)  
10 10 to a final concentration of about 100 ng/ml. About one hundred microliters of this diluted streptavidin were  
25 then added to each well. Each plate was covered and incubated for about 1 hour at 4C with mixing. Each plate was washed four times with wash buffer at 4C  
30 15 after which about 100 microliters of Enhancement Solution (Wallac; item number 1244-105) was added to each well. The plate was then mixed for at least 5  
35 minutes at room temperature.

40 Delfia time resolved fluorescence was read  
20 with a Wallac Victor 1420 multilabel counter using an excitation wavelength of 340 nm and an emission wavelength of 613 nm.

45 B. Assay for the 42 Amino Acid Form of A-Beta

25 A rabbit polyclonal antibody which specifically recognizes residue 42 of A-beta (Quality Controlled Biochemicals, MA 01748-2215; item number  
50

- 86 -

44344) was diluted to about 2.5 micrograms/ml in the  
diluent Superblock TBS (Pierce Chemical Co. Rockford,  
IL 61105) and was added to each Reacti-Bind goat-anti-  
rabbit IgG-Fc specific plate (Pierce Chemical Co.;  
Rockford, IL 61105 item number NC1513). The plates  
were then incubated for about one hour at room  
temperature on a plate mixer with vigorous mixing.  
Following incubation, the plates were washed with wash  
buffer TBS-Tween (0.5 percent) four times, and the non-  
specific binding sites were blocked with about 100  
microliters per well of Superblock TBS.

The standard curve samples were prepared from  
a stock solution of A-beta-1-42 peptide in DMSO at a  
concentration of 5 micrograms/ml. The highest standard  
was prepared at a final concentration of 40,000 pg/ml  
in Superblock TBS. Sequential dilutions were made  
using Superblock TBS to obtain final concentrations of  
20,000, 10,000, 5,000, 2,500, 1,250, 625, 312.5,  
156.25, 78.125 and 39.06 pg/ml.

About one hundred microliters per well of  
each standard, or 17.5 microliters per well of each  
sample was applied to each well of the microtiter  
plates to which about 100 microliters of Superblock TBS  
had been previously added. The plates were covered and  
incubated at about 4C overnight with mixing. The  
plates were then washed four times with wash buffer  
containing about 0.5 percent TBS-Tween-20™ at 4C.

5

- 87 -

10

15

20

25

30

35

40

45

50

55

Detection antibody, m4G8-biotin (Senetek; St. Louis, MO; item number 240-10) was diluted to about 0.5 micrograms per ml in a solution containing Superblock TBS, 2 percent normal mouse serum, 2 percent normal goat serum, and about 100 microliters of the diluted antibody were added to each well. Each plate was covered and incubated for about 2 hours at 4C with mixing. Each plate was then washed four times with standard wash buffer at 4C.

10 Europium labeled streptavidin (Wallac, Gaithersburg, MD; item number 1244-360) was diluted 1:1000 in Assay Buffer (Wallac, Gaithersburg, MD; item number 1244-106) to a final concentration of about 100 ng/ml. About one hundred  $\mu$ l per well were then added to each well. The plates were covered and incubated for about 1 hour at 4C with mixing. Each plate was washed four times with wash buffer at 4C, after which about 100  $\mu$ l of Enhancement Solution (Wallac, Gaithersburg, MD; item number 1244-105) were added to each well. Each plate was then mixed for at least 5 minutes at room temperature.

Delfia time resolved fluorescence was read with a Wallac Victor 1420 multilabel counter using an excitation wavelength of about 340 nm and an emission wavelength of about 613 nm.

- 88 -

Example IV: Isolation of a Beta-Secretase cDNA

Approximately 864,000 independent clones were screened in the assays described above. Positive clones were selected based on the following two criteria:

- 1) increased A-beta 42 and/or total A-beta levels by more than two standard deviation points as compared with the plate average; and
- 2) selective skewing of the A-beta 42/ total A-beta ratio towards A-beta 42 production.

Based on these criteria, about 144 putative positive pools of cDNAs were replated onto new master plates for a second round of analysis. An aliquot of the cDNA from each pool was transfected into 293T-APPsw101 (described above) using procedures described above. Transfections were conducted in duplicate. Those pools of cDNAs which consistently met positive clone criteria were subsequently replated and diluted from pools of about 100 clones to pools of about 20 clones and then to single clones. Single clones were diluted 1:10 into pCMV1 vector DNA since the assumption was made that severe over-expression of a protease might result in toxicity to the cells.

One pool, termed A-11, exhibited a dramatic skewing of the A-beta 42/A-beta total ratio. In

- 89 -

duplicate assays, this pool gave a ratio of about 0.8, and 1.0, respectively, whereas the plate average ratio for all plates ranged from about 0.5 to about 0.7. When A-11 was subsequently re-plated, it showed a modest yet consistent skewing of the A-beta 42/A-beta total ratio. In addition, A-11 had a standard deviation of greater than 2 for the increase in A-beta 42 levels over the plate average. The ratio skewing seemed to be somewhat dependent upon the density of cells at the time of transfection. When the cells were less dense at time of transfection, the skewing appeared to be more robust. As pool A-11 was broken down, it exhibited enhanced signal strength by both of the above criteria.

The results of the three assays for pool A-11 as well as the plate average are set forth below in Table II as total Europium counts. The data are presented as an average of duplicate transfections where each transfection was run in duplicate assays.

- 90 -

Table II

Assay	Pool A-11	Plate Average
Re-Arrayed Pool of 100		
A-beta 42	94,397	72,858
Total	177,356	164,334
A-beta 42/Total	0.53	0.44
Pool of 20		
A-beta 42	178,212	133,402
Total	448,660	388,093
A-beta 42/Total	0.397	0.34
Single Clone ( $\beta$ -Secretase)		
A-beta 42	327,505	230,518
Total	295,438	363,141
A-beta 42/Total	1.1	0.63

When pool A-11 was diluted to single clones, several of these clones were sequenced using standard methods. All of the sequenced clones possessed an identical nucleotide sequence. After sequencing, one of these clones was inserted into the vector pCMVi-beta-secretase. Further sequence analysis indicated that the sequence possessed many sequence

- 91 -

similarities to the aspartic protease family, and encoded a novel aspartic protease containing both a putative signal sequence and a transmembrane domain. The cDNA sequence of this aspartic protease, identified as a novel sequence encoding beta-secretase, is set forth in Figure 1A and 1B. The putative amino acid sequence as translated from the cDNA is set forth in Figure 4, and this sequence possesses certain motifs characteristic of an aspartic proteases, such as two putative active site domains. The propeptide consists of amino acids 23-45. The signal peptide spans amino acids 1-22, and the transmembrane domain spans amino acids 461-477. The enzyme appears to be cleaved *in vivo* between amino acids 22 and 23 in some cases to remove the signal peptide, and in addition between amino acids 45 and 46 to remove the propeptide in other cases. The active site domains are believed to be amino acids 93-96 and 289-292. Hence, it is believed that the mature, active beta secretase enzyme spans amino acids 46-501. Based on computerized three dimensional structure analysis of human beta amino acid sequence, it is expected that the fragments 62-420, 73-420, 83-420, 90-420, 62-417, 73-417, 83-417, 90-417, 62-410, 73-410, 83-410, 90-410, 62-402, 73-402, 83-402, and 90-402 would all be biologically active.

- 92 -

Example V: Identification of the Mouse and Rat  
Orthologs of Beta-Secretase

To identify the mouse and rat beta-secretase cDNAs, a FASTA search was performed using the full length human beta-secretase cDNA to search an Amgen database. For conducting this search, the scoring matrix used was GenRunData.fastadna.cmp, together with a constant pam factor. The Gap creation penalty and Gap extension penalty were set for 12.0 and 4.0, respectively. Four ESTs were identified, three for mouse and one for rat.

The three mouse sequences, which were 351 base pairs, 411 base pairs, and 364 base pairs in length were found to overlap. Thus, a complete mouse beta-secretase cDNA was obtained. This mouse cDNA was about 1923 base pairs in length and included about 95 nucleotides of 5 prime untranslated sequence, about 1503 nucleotides of open reading frame encoding 501 amino acids, and approximately 325 nucleotides of 3 prime untranslated sequence.

The single rat clone obtained encoded full length beta-secretase and was about 2158 nucleotides in length and contained 427 nucleotides of 5 prime untranslated sequence and 225 nucleotides of 3 prime untranslated sequence. The rat beta secretase is also 501 amino acids in length.



- 93 -

The full length mouse and rat beta-secretase cDNAs share approximately 93 percent and 91 percent identity, respectively, at the nucleic acid level with the human full length beta-secretase cDNA.

The mouse cDNA sequence containing only the full length coding region is set forth in Figure 2A and 2B (SEQ ID NO:2) and the corresponding amino acid sequence is set forth in Figure 5 (SEQ ID NO:5).

The rat cDNA sequence containing only the full length coding region is set forth in Figure 3A and 3B (SEQ ID NO:3) and the corresponding amino acid sequence is set forth in Figure 6 (SEQ ID NO:6).

Example VI: Northern Blot Analysis of Beta-Secretase Transcripts

Northern blot analysis was performed to identify those tissues in which the beta-secretase transcript is present. A probe for use in Northern blot analysis was generated by digesting the human  $\beta$ -secretase cDNA with *Pst* I for about three hours at 37°C and running the restriction digest on an 0.8 percent agarose gel to separate the fragments. The approximately 772 base pair ("bp") *Pst* I fragment extending from nucleotide 318 to nucleotide 1090 of the cDNA was isolated and gel purified using the QiaQuick®

- 94 -

gel purification system (Qiagen, Chatsworth, CA). The isolated, gel pure fragment was quantitated by estimation on a one percent agarose gel. About 25 ng of this fragment was denatured by boiling for 5 minutes, and then quenching on ice for 2 minutes. The fragment was then radioactively labeled with alpha 32P-dCTP using the High Prime DNA labeling kit (Boehringer Mannheim, Indianapolis, IN) according to the manufacturer's protocol. Human multiple tissue northern blots were purchased (Clontech, Palo Alto, CA) and first prehybridized in Clontech Express™ hybridization buffer for about one hour at about 65°C. Following prehybridization, the labeled probe was denatured by boiling for about five minutes and quenching on ice for 2 minutes, and then added to the hybridization buffer containing the Northern blots. The blots were allowed to hybridize for about two hours at about 65°C. After hybridization the blots were washed in 2xSSC for 30 minutes at room temperature, followed by 3 successive washes in 0.2xSSC containing 0.1 percent SDS at about 60°C for 30 minutes. The blots were dried briefly and exposed to autoradiography film for 72 hours at about -80°C.

The results are shown in Figure 7. The lane contents are described in the Brief Description of the Figures. Three different RNA transcripts of approximately 7 kb, 4.4 kb and 2.6 kb are apparent.

- 95 -

All are of low abundance and expression is detected in most tissues. The highest levels are observed in pancreas and brain.

Example VII: Beta-Secretase Protein Detection

An antibody to the carboxy terminus of beta-secretase was raised as follows. A synthetic peptide with the sequence CLRQQHDDFADDISLLK (SEQ ID NO:7) corresponding to amino acids 485-501 of beta-secretase was generated using standard methods (see below). The peptide was then conjugated to the carrier protein KLH by adding 5 mg of peptide to Pierce maleimide conjugation buffer (Pierce Chemical Co.; Rockford, IL; item number 77164), after which 0.5 ml of Pierce maleimide activated KLH (5 mg) (Pierce Chemical Co.; item number 77105) was added. The solution was incubated for about 2 hours at room temperature, and then run through a D-Salt™ Dextran column (Pierce Chemical Co.; item number C43233) to remove the EDTA. A one ml fraction from the column was mixed 1:1 with Titermax Research Adjuvant (CytRx Corp., Norcross, Georgia; item number R-10), prepared as an oil and water emulsion using 18 gauge double-hubbed needles (Popper and Sons, New Hyde Park, N.Y) and loaded into 1 c.c. syringes fitted with 21 gauge needles. Three New

5

- 96 -

10

15

20

25

30

35

40

45

50

Zealand White rabbits were each injected intra-muscularly (IM) at two injection sites (0.05 ml per site). Four and six weeks later, the rabbits were boosted IM, again at two sites, 0.05 ml per site. The first sample bleed of 5 ml was drawn at 6 weeks. Two weeks later the second test bleed was drawn and tested for immunoprecipitation of  $\beta$ -secretase from transfected cells. The results indicated that a beta-secretase specific antibody had been generated.

10

The antibody was coupled to Protein A Sepharose using standard procedures for use in immunoprecipitation assays. The preimmune serum (negative control) was coupled in the same manner. Coupling was conducted as follows:

15

Protein A Sepharose CL-4B beads (Amersham Pharmacia Biotech, Inc, Piscataway, NJ; item number 17-0780-01) were suspended at about 250 mg/ml in BSA/TBS (50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 2% BSA). About 1 ml of the resuspended beads was mixed with about 250  $\mu$ l of antiserum. The mixture was incubated at room temperature, rocking gently, for about one hour. The beads were then washed with about 10 ml of borate buffer (0.2 M sodium borate, pH 9.0). After about five minutes of gentle rocking, the mixture was spun at 3000g for about five minutes and the supernatant was discarded. This washing procedure was then repeated, after which the beads were resuspended in about ten ml

55

- 97 -

of borate buffer. Before adding the coupling agent,  
about 100  $\mu$ l of the slurry was removed for gel  
electrophoresis analysis (tube A, see below).

The antiserum was then covalently coupled to  
the beads by adding about 50 mg of solid dimethyl  
pimelimidate (DMP, Pierce Chemical Co., Rockford, IL;  
item number 21666). The final concentration of DMP was  
about 20 mM and had a pH of at least 8.3. After  
incubating the mixture at room temperature for about  
thirty minutes with gentle rocking, a 100 microliter  
aliquot was removed (tube B, see below); the remainder  
of the mixture was spun at about 3000 x g for about 5  
minutes.

The coupling reaction was terminated by  
washing the beads in about 10 ml of 0.2M ethanolamine,  
pH 8.0. After mixing, the solution was spun at about  
3000 x g for about 5 minutes. The beads were  
resuspended in about 10 ml of ethanolamine and  
incubated for about 2 hours at room temperature with  
gentle rocking. The beads were then spun at about 3000  
x g for about 5 minutes, after which the pellet was  
washed briefly with about 2 ml glycine (100 mM glycine,  
pH 3.0), followed by spinning again at about 3000 x g  
for about 5 minutes.

The beads were rinsed with about 2 ml Tris  
(100 mM, pH 8.0) and spun at about 3000 x g for about 5  
minutes, after which the beads were resuspended in

- 98 -

about 1 ml PBS containing 0.01 percent thimerosal.

The beads were stored at 4°C. An aliquot (about 20 µl)

of this final product (tube C) was run on SDS-PAGE gel along with tubes A and B (see above) to check the

efficiency of coupling. The beads were prepared for SDS-PAGE by pelleting and resuspending them in about 30

µl of SDS sample buffer (Novex, San Diego, CA; item

number LC2676), after which they were heated at about

85°C for about 10 minutes and then run on a 10 percent

Tris Glycine gel (Novex, San Diego, CA) at 100V for

about 2-3 hours. The gel was and stained with

Coomassie blue.

To evaluate the presence of beta-secretase in human brain tissue, human Alzheimer's disease and age-

matched control brains were obtained from Sun Health Research Institute (Sun City, Arizona) and homogenized according to the following protocol:

Pieces of parietal cortex of approximately 1 gram wet weight were finely chopped while frozen, and

then placed in a Dounce glass homogenizer. Lysis

buffer (50 mM Tris, pH 7.5, 150 mM NaCl, 1 percent

NP-40, 12 mM CHAPS, 0.2 percent BSA, 2 mM EDTA, 20 mM

PMSF, 10 µM leupeptin, 1 µM pepstatin A, 2 µg/ml

aprotinin, 0.1 mM Pefabloc) was added in a proportion

of 2 ml buffer per 1 g brain tissue, and the mixture

was homogenized using 50 strokes with a tight fitting

- 99 -

plunger. The homogenate was spun at 20,000 x g for about 10 minutes at 4°C. The supernatant was used for the immunoprecipitation and Western blotting as described below. Remaining material was stored at -80°C. The supernatant concentration was assumed to be approximately 500 mg protein /ml based on the starting wet weight.

The supernatants were immunoprecipitated as described immediately below with either the antibody or with preimmune serum coupled to protein A Sepharose (prepared as described above), separated on SDS-PAGE, and Western blotted with the pooled antisera. As a positive control, lysates made from 239T-APPsw101 cells described above transiently transfected with the full length beta-secretase expression construct were analyzed under the same conditions.

For each immunoprecipitation, about 50 µl of brain supernatant prepared as described above (about 25 mg wet weight equivalent) or the lysate made from 101 cells transiently transfected with the beta-secretase expression construct (about 1/20 of the volume of each dish) were used. The brain supernatant and the lysate were precleared in bulk (about 100 µl homogenate containing about 25 µl Protein A Sepharose) by incubating at about 4°C for 30 minutes with gentle rocking. The tubes were centrifuged at about 960 x g

- 100 -

for about 5 minutes. The supernatants (about 50  $\mu$ l) were immunoprecipitated with about 40  $\mu$ l of the Protein-A- beta-secretase antisera or Protein-A-preimmune serum. The volume of the immunoprecipitation was brought up to about 500  $\mu$ l with TBS (50 mM Tris, pH 7.5, 150 mM NaCl) and the reaction was carried out at about 4°C for about 3 hours with rocking. The beads were pelleted at about 960 x g for about 5 minutes and washed three times with STEN buffer (1x STEN = 50 mM Tris-HCl, pH 7.5, 150 mM NaCl, 2 mM EDTA, 0.2 percent NP-40) for about 15 minutes at about 4°C with rocking. The beads were pelleted after each wash at about 960 x g for about 5 minutes. The first wash buffer was 0.5 M STEN (1x STEN + 0.5 M sodium chloride, 5  $\mu$ g/ml leupeptin, 5  $\mu$ g/ml aprotinin). The second wash buffer was SDS-STEN (1x STEN + 0.1% SDS, 5  $\mu$ g/ml leupeptin, 5  $\mu$ g/ml aprotinin), and the third wash buffer was 1x STEN. After washing, about 15  $\mu$ l of SDS-Sample Buffer (Novex, San Diego, CA; item number LC2676) was added and the samples were stored at -20°C overnight.

The samples were heated at about 95°C for about 5 minutes and loaded onto a 10-20 percent Tris-Tricine gel (Novex San Diego, CA ; item number EC6625). The gel was run at 100V for 2-3 hours and then transferred to PVDF (Novex, San Diego, CA; item number



- 101 -

LC2002) using the Bio Rad Trans Blot Cell at about 400  
mA for about 2 hours at 4°C in a solution of 200 mM  
glycine, 25 mM Trizma base and 20 percent methanol.

The Western blot was then analyzed for the  
presence of beta secretase as follows. The Western blot  
was wetted in methanol for about 30 seconds then rinsed  
with TBST (10 mM Tris-HCl, pH 8.0, 150 mM sodium  
chloride, 0.05 percent Tween-20) before blocking with 5  
percent dry milk in TBST for about one and one-half  
hours at room temperature, with rocking. The Western  
blot was incubated with primary beta-secretase antibody  
(prepared as described above) that had been diluted  
about 1:1000 in PBS containing 1 percent BSA and 0.1  
percent Tween-20. Incubation was overnight at about  
4°C with rocking. After incubation, the Western blot  
was washed three times in TBST for about 5 minutes per  
wash with rocking at room temperature.

The secondary antibody used to detect beta-  
secretase antibody was alkaline phosphatase conjugated  
Goat Anti-rabbit IgG (Promega, Madison, WI; item number  
S373B). This antibody was diluted at about 1:7500 in  
alkaline phosphatase buffer (100 mM Tris-HCl, pH 9.2,  
100 mM sodium chloride, 5 mM MgCl<sub>2</sub>) and added to the  
Western blot. After about 30 minutes of incubation at  
room temperature, the Western blot was washed as  
described above and developed about 5-10 minutes with  
BCIP/NBT (Kirkegaard and Perry Labs, Gaithersburg, MD;

- 102 -

item number 50-81-08). To stop development, the Western blot was rinsed in stop solution (20 mM Tris-HCl, pH 8.0, 5 mM EDTA) and dried.

The results are shown in Figure 8. "P" represents immunoprecipitations with preimmune serum, and "A" represents immunoprecipitations with anti-beta secretase antibody. "Cells" refers to human cultured cells transfected with the human beta-secretase cDNA; "AD" refers to brain tissue homogenate from Alzheimer's disease patients; and "Control" refers to brain tissue homogenate from normal people. As can be seen, a major band of approximately 70 kilo daltons is apparent in lane 2. A band of approximately the same molecular weight is present in lanes 4 and 6 as well. This result indicates that the transfected cells are expressing the beta secretase protein which is observed in human brain. The molecular weight of beta-secretase calculated from the amino acid sequence is about 55.8 kilo daltons, suggesting that the protein expressed in cells and brain could be glycosylated.

Example VIII: Preparation of a Human Beta-Secretase Mutant

A hemagglutinin (HA) epitope tag was added in frame to the C-terminus of the full length beta-

- 103 -

secretase cDNA for the purpose of detection with the monoclonal anti-HA antibody HA.11 (obtained from BabCo, Berkeley CA). The HA epitope tag consists of 11 amino acids and was used as a marker tag, enabling the surveillance of the beta-secretase-HA fusion polypeptide via anti-HA antibody detection.

The C-terminal HA tagged form of beta-secretase was generated by PCR in two steps. For the first step, the forward primer had 5 prime to 3 prime sequence corresponding to nucleotides 1047-1070 of the beta-secretase cDNA and had the sequence:

TGACTCTCTGGTAAAGCAGACCCA (SEQ ID NO:8)

The reverse primer for this first round of PCR corresponded to nucleotides 1938-1957 of beta-secretase together with the first 18 base pairs of the HA tag and had the sequence:

AGGCACGTCGTAAGGGTACTTCAGCAGGAGATGTCA (SEQ ID NO:9)

The PCR conditions for this reaction were: 97C for 60 seconds; 97C for 30 seconds, 60C for 30 seconds and 72C for 1 minute for twenty five cycles.

This round of PCR generated a 928 base pair product. This PCR product was then purified using the Wizard<sup>™</sup> PCR Preps DNA Purification System (Promega, Madison,

- 104 -

WI). This PCR product was then used as template for the second PCR reaction. This reaction used the identical forward primer and a reverse primer corresponding to all 11 amino acids of the HA tag followed immediately by a termination codon and a *BglII* restriction site. This primer had the sequence:

TGAAGATCTTCATCCGCTGGCATAATCAGGCACGTCGTAAGGGTA (SEQ ID NO:10)

PCR conditions for this reaction were identical to those used for the first reaction. The 955 base pair PCR product that resulted from this second round of PCR was purified as described above and digested with *BsrGI* and *BglII* overnight at 37C. The digested fragment was gel purified from a 0.8 percent agarose gel using the QiaQuick gel purification system (Qiagen, Chatsworth, CA) and subsequently subcloned into *BsrGI/BglII* digested pCMVi-beta-secretase.

A mutant DNA construct encoding D93A human beta secretase was prepared via PCR. Each fragment contained the D93A mutation and 24 base pairs of overlapping sequence. The first or forward fragment contained the D93A mutation in the sense orientation at the extreme 3 prime end of the fragment, while the second or reverse fragment contained the D93A mutation at the extreme 5 prime end of the fragment in the

- 105 -

antisense orientation. Next, these PCR products were purified and combined together (at equimolar ratio) in a second amplification reaction using just the two outer primers for amplification, thus giving rise to a full length double stranded 953 bp *Bam*HI-*Bsr*GI 008 fragment of beta secretase containing the D93A mutation. This PCR product was then subsequently used to replace the *Bam*HI-*Bsr*GI fragment of the native beta-secretase cDNA.

The forward first step PCR product was generated using a 5 prime forward primer which is identical to bases 211-233 of the beta-secretase cDNA sequence. This primer has the sequence:

GTGCCGATGTAGCGGGCTCCGGA (SEQ ID NO:11)

A 3 prime reverse primer was generated which corresponds to bases 720-743 of human beta-secretase cDNA except that the T at position 731 was changed to G thus changing the aspartic acid residue at position 93 of the peptide sequence to an alanine residue (on the antisense strand). This primer has the sequence:

CTGCTGCCTGTAGCCACCAGGATG (SEQ ID NO:12)

The reverse first step PCR product was generated using a 5 prime forward primer in which the A

- 106 -

at base 731 was changed to a C, thereby creating a construct that encoded an alanine at position 93 of the human beta-secretase polypeptide sequence (on the sense strand). This primer has the sequence:

CATCCTGGTGGCTACAGGCAGCAG (SEQ ID NO:13)

The 3 prime reverse primer used in this PCR reaction is identical to bases 1230-1254 of the beta secretase human cDNA. This primer has the sequence:

CACCGGCACAATGATCACCTCATAA (SEQ ID NO:14)

PCR reactions for the forward first step reaction were conducted in an MJ Research PTC 200 thermal cycler (MJ Research, Watertown, MA) using the following conditions: 97C for 90 seconds; 98C for 30 seconds, 60C for 30 seconds and 72C for 1 minute for twenty five cycles. The two PCR products were then purified using the Wizard™ PCR Preps DNA Purification System (Promega, Madison, WI).

For the second step PCR reaction, the two PCR fragments generated above were added in an approximately equimolar ratio, and were subsequently amplified using the following primers:

- 107 -

GTGCCGATGTAGCGGGCTCCGGA (SEQ ID NO:15)

CACCCGCACAATGATCACCTCATAA (SEQ ID NO:16)

The PCR conditions for this reaction were identical to those used in the first step reactions. The PCR products were purified using the Wizard™ PCR Preps DNA Purification System (Promega, Madison, WI), restriction digested with *Bam*HI and *Bsr*G,I and gel purified on a 0.8 percent agarose gel using the QiaQuick gel purification system (Qiagen, Chatsworth, CA). This fragment was then subcloned into the *Bam*HI/*Bsr*GI digested pCMVi-beta-secretase/ HA tag construct.

In a separate procedure, the human APP-695 cDNA was subcloned into the CMV based expression vector pCMVi (Cell & Molecular Technologies Inc., LaValllette, New Jersey) as a *Hind*III- *Not*I fragment to generate the vector pCMVi-APPwt. A derivative of 293 cells containing the SV40 large T antigen (Cell & Molecular Technologies, LaValllette, NJ ) maintained in high glucose Dulbecco's modified Eagle's medium (DMEM) supplemented with 10 percent fetal calf serum was co-transfected with pCMVi-APPwt and pRSV-Puro by the calcium phosphate method as described (Gorman et al., supra). Transfected cells were cultured in medium containing puromycin at a concentration of about 5

- 108 -

micrograms/ml . Puromycin resistant colonies were selected about 48 hours after transfection. Fourteen days later, individual resistant colonies were isolated, expanded, and analyzed by Western dot blot for APP expression using the monoclonal antibody 22C11 (Boehringer Mannheim Corp.) which recognizes secreted APP.

One positive colony was expanded to generate a cell line referred to as "293T/APPwt".

To assess the activity of the D93A mutant, 293T/APPwt cells were transfected with beta-secretase cDNA (either wild type or D93A mutant) or with control plasmid (no beta-secretase insert) as follows.

The 293T/APPwt cells were plated out at a density of about  $2.8 \times 10^5$  cells per well in a 6-well tissue culture plate (Falcon) and grown for 2 days at 37°C in an atmosphere of about 5 percent CO<sub>2</sub>. The cells were transfected the morning of day 3. Prior to transfection, all plasmids to be used were resuspended in water. Two polystyrene tubes were set up for each well. One tube received about 3 µg total DNA (control plasmid was diluted 1:10 as compared with the plasmids containing beta-secretase inserts) and 1 ml of Opti-MEM I reduced serum media (Gibco-BRL, Grand Island, New York). The second tube received about 14 µl DMRIE-C (Gibco-BRL) and 1 ml Opti-MEM I. The tubes were then mixed together and incubated at room temperature for



- 109 -

about 30 minutes. The media was then removed from the cells and the DNA/ DMRIE-C/ Opti-MEM I mixture (2 ml total volume) was added to each well. The cells were incubated at about 37°C in about 5 percent CO<sub>2</sub> for about 5 hours. Following the incubation, about 2 ml of Opti-MEM I containing 20 percent fetal bovine serum (Gibco-BRL) were added to each well of cells, after which the cells were incubated at about 37°C, in an atmosphere of about 5 percent CO<sub>2</sub> overnight. The following morning (day 4) the media was removed and 2 ml of fresh DMEM containing 10 percent fetal bovine serum (Gibco-BRL) were added, and the cells were then incubated at about 37°C in an atmosphere of about 5 percent CO<sub>2</sub> overnight. The following day (day 5) the conditioned media was collected, and sodium dodecyl sulfate (Sigma, St. Louis, MO) was added to a final concentration of about 0.1 percent. The samples were stored at about -80°C.

Figure 9A and 9B show the results of the transfection. Conditioned media of cells transfected with the wild type beta-secretase expression plasmid contain significantly higher levels of A-beta 42 (Fig. 9A) and A-beta 40 (Fig. 9B) than conditioned media of cells transfected with either the control plasmid or conditioned media of cells transfected with the beta-secretase D93A-HA mutant. This suggests that D93 is part of the active site of human beta-secretase.

5

- 110 -

10

Example IX: Purification of Soluble Beta Secretase-Fc  
Fusion Polypeptide

15

A human IgG1 Fc gene (the sequence of which  
5 is set forth in Genbank as accession number X70421) was  
amplified by polymerase chain reaction and subcloned  
into the vector pBluescript SK- (Stratagene, La Jolla,  
CA). PCR was conducted using a MJ Research PTC-200  
20 thermal cycler for 25 cycles with the following  
parameters: 94C for 1 minute; 55C for 1 minute; and 72C  
10 for 1 minute. The polymerase used was Pfu polymerase  
(Stratagene, La Jolla, CA). The following primers were  
used as forward and reverse primers, respectively for  
this PCR reaction:

30

15

CGGGATCCGGTCACCGACAAAACCTCACACA (SEQ ID NO:20)

35

GCTCTAGAAGCTTCTGCAGGTCGACTCATTTACCCGGAGA (SEQ ID NO:21)

40

20 The forward primer incorporated 5' *Bam*HI and  
*Bst*EII sites at the 5' end of the PCR product, and the  
reverse primer incorporated *Sal*I, *Pst*I, *Hind*III, and  
*Xba*I sites at the 3' end of the 723 base pair PCR  
45 fragment that was generated using these primers. The  
PCR fragment was purified using Qiaquick™ purification  
25 kit (Qiagen, Chatsworth, CA), and digested with *Bam*HI  
and *Xba*I. After a second round of gel purification,

50

55

- 111 -

the PCR fragment was ligated into the vector  
pBluescript SK- (Stratagene, La Jolla, CA) that was  
previously digested with *Bam*HI and *Xba*I.

An 1195 base pair fragment of the human beta-  
secretase cDNA (nucleotides 276 to 1471) was excised  
from the full length cDNA by digestion with *Eco*RI and  
*Bsr*GI and gel purified. This fragment contained the 5'  
untranslated region and the coding region up to amino  
acid 245 of the human beta secretase cDNA. A second  
fragment of human beta-secretase cDNA encoding amino  
acids 246 to 460 (ending exactly at the start of the  
putative transmembrane domain) was amplified by PCR  
using procedures described above for the IgG1 fusion  
gene. Amplification forward and reverse primers for  
this reaction were as follows:

CGACCACTCGCTGTACACAGGCAG (SEQ ID NO:22)

GTCGGTGACCGCATAGGCTATGGTCATGAGGGT (SEQ ID NO:23)

The forward primer incorporated the  
endogenous 5' *Bsr*GI site, and the reverse primer  
incorporated a 3' *Bst*EII site for in-frame ligation of  
the PCR product into the IgG1 fusion gene cassette.  
The resulting 672 base pair PCR fragment was purified  
using the Qiaquick<sup>TM</sup> purification kit (Qiagen,

- 112 -

Chatsworth, CA), digested with *BsrGI* and *BstEII*, and gel-purified.

Separately, the human IgG1 fusion cassette was digested with *EcoRI* and *BstEII* and gel-purified. A 3-way ligation was then conducted by combining the 1195 base pair *EcoRI/BsrGI* human beta-secretase fragment and the 672 base pair *BsrGI/BstEII* beta secretase PCR fragment with the *EcoRI/BstEII* human IgG1 fusion cassette. The completed beta secretase fusion gene was called beta-secretase-Fc and contained 3 additional amino acids (alanine-valine-threonine) at the junction between the beta-secretase and IgG1 coding regions.

The beta-secretase-Fc fusion gene was excised from the plasmid pBluescript SK- by digestion with *XhoI* and *NotI*, and gel-purified. The mammalian expression vector pCMVi (Cell & Molecular Technologies Inc. Lavallette, NJ) was also digested with the same enzymes and gel-purified, after which the beta-secretase-Fc fusion gene fragment was ligated into the *XhoI/NotI* digested pCMVi vector.

Human embryonic kidney 293T cells (Cell & Molecular Technologies Inc., Lavallette, NJ) stably transfected with the beta-secretase-Fc fusion construct by the standard calcium phosphate method (Corman et al, *DNA Prot. Eng. Tech.* 2: 3-10 [1990]) were conditioned for either four or five days in Opti-Mem reduced serum medium (GibcoBRL, Grand Island, NY; item number 31985).

- 113 -

After filtration with a 0.45  $\mu$ m filter, conditioned media was concentrated and buffer exchanged to column equilibration buffer (PBS containing 2mM Chaps). The concentrated conditioned medium was then loaded onto a one milliliter recombinant protein A column (Amersham Pharmacia Biotech, Inc, Piscataway, NJ; item number 17-5079-02) which had been pre-equilibrated with 10 column volumes of PBS containing 2mM Chaps™ at about 4C. After loading the sample on the column, the column was washed with 10 column volumes of column equilibration buffer, and the beta-secretase-Fc fusion protein was eluted from the column with a step gradient to 100 mM Sodium Citrate, pH 2.8 containing 2mM Chaps™. One milliliter fractions were collected. Before collection, fraction collection tubes were loaded with 300 ml of 1M Tris base, pH 9.2, to neutralize fractions upon elution. Final pH of each 1.3 ml fraction was approximately pH 7.5. Fractions containing beta-secretase-Fc fusion polypeptide were pooled, concentrated, and buffer exchanged to TBS containing 2 mM Chaps for storage.

Figure 10 shows a Coomassie stained SDS gel of the purified beta-secretase-Fc fusion protein (lane 2) alongside molecular weight markers (lane 1). Only the beta-secretase-Fc fusion polypeptide and some cleaved Fc polypeptide were detectable. The predicted

- 114 -

molecular weight of the fusion protein is 75.8 kilo daltons. The higher observed molecular weight is presumably due to glycosylation.

5 Example X: Beta-Secretase Activity Assay

Recombinant native as well as mutant beta secretase polypeptides were assayed for activity by measuring the ability of the molecule to cleave various A-beta peptides.

All peptides were synthesized by the Fmoc (fluorenylmethoxycarbonyl) / t-butyl based solid phase peptide chemistry method using standard procedures. An ABI 431A peptide synthesizer (Perkin Elmer Corp., Foster City, CA) was used with a single coupling program to carry out the chain assembly. Commercially available preloaded Fmoc-AAA-HMP derivatized polystyrene resin (Midwest Biotech, Fishers, IN or Calbiochem, San Diego, CA) was used to prepare the C-terminal amino acid. Subsequent amino acids were coupled in 20 fold excess as HOBT (hydroxybenztriazole) esters using carbodiimide activation. The side-chain protecting groups for each amino acid were as follows: Arg(Pbf; 2,2,4,6,7-pentamethyldihydrobenzofuran-5-sulfonyl), Asn(Trt; trityl), Asp(OtBu; O-tert-butyl), Cys(Trt; trityl), Cys(Acm; acetamidomethyl), Gln(Trt; trityl), Glu(OtBu; O-tert-butyl), His(Trt; trityl).

- 115 -

Lys(Boc; tert-butoxycarbonyl), Ser(tBu; tert-butyl),  
Thr(tBu), and Tyr(tBu; tert-butyl). Upon removal of  
the final N-terminal Fmoc with 20 percent piperidine in  
N-methylpyrrolidone, side-chain protecting groups were  
then removed and the peptide(s) were cleaved from the  
resin by treatment with TFA (trifluoroacetic acid):  
triisopropylsilane: water (92.5, 2.5, 5 v/v) for about  
4 hours. The resulting suspension was filtered, and  
the filtrate volume reduced by roto-evaporation. The  
crude peptides were precipitated and washed with ether,  
followed by drying in-vacuo.

The linear (fully reduced) peptide  
intermediates were purified by HPLC prior to either an  
equilibrium oxidative refold or a two step oxidative  
cyclization process. The connectivity of those  
peptides containing one or two disulfides is  
unambiguous as a result of using orthogonal cysteine  
protection and oxidative-cyclization methods as  
follows. The first disulfide bond was formed using air  
oxidation (Cys(Trt) protection) and the second  
disulfide bond was formed by iodine treatment on the  
bis-Cys(Acm) containing -cyclic peptide intermediate.  
For the two peptides containing three disulfide  
bridges, oxidation was carried out using 10 percent  
DMSO (dimethylsulfoxide) oxidation under acidic  
conditions.

- 116 -

Each crude peptide was dissolved in 8M guanidine containing 100 mM DTT (dithiothreitol) and purified to at least 95 percent homogeneity by preparative reverse-phase HPLC using a Vydac C18 (2.5 cm x 25 cm) column (Vydac Corp., Hesperia, CA) with a linear gradient of 0.1 percent TFA (v/v) in water and 0.05 percent TFA (v/v) in acetonitrile.

The composition of each peptide was assessed using electro-spray ionization (ESI) mass spectrometry and amino acid analysis. Mass spectra for each synthetic peptide was obtained on a Sciex API (Perkin Elmer Corp., Foster City, CA) single quadropole mass spectrometer, and reported as  $m/z$  (M+1). All mass spectral samples were obtained as fractions off of the preparative HPLC purification.

Amino acid analyses of each peptide were performed on an ABI 420A hydrolyzer/derivatizer (ABI, Foster City, CA) using a 130A separation system (ABI, Foster City, CA). The peptides were hydrolyzed using 6N HCl at 200° for 30 minutes and then derivatized using the defined instrument protocol in the ABI AAA420A Operator's Manual as PTC (phenylisothiocynate) derivatives. The amino acid mixture was then separated by HPLC on a Brownlee PTC C18 column (ABI, Foster City, CA), 5 micron pore size, 2.1 x 220 mm, with a linear gradient of water and acetonitrile. Both solvents were



- 117 -

buffered with sodium acetate to a pH of about 5.4. The amino acid composition of each peptide was then determined by comparison of the unknown peak ratios with an equimolar amino acid standard. Each peptide generated experimental data that conformed with expected theoretical values.

The sequence of the synthesized peptides (from amino to carboxy terminus) is set forth in Table III below:

Table III

Peptide	Peptide Sequence	
Sw	EVNLDAEF	SEQ ID NO:17
WT	EVKMDAEF	SEQ ID NO:18
MV	EVKVDAEF	SEQ ID NO:19

To facilitate analysis of the assays, all of the peptides were labeled with dinitrophenol (DNP) at the amino terminus.

DNP was incorporated during synthesis of the peptide substrates as dnp-glutamic acid.

Each assay was conducted as follows. The purified beta secretase-Fc (amounts indicated in Fig. 11) was combined separately with each peptide substrate

- 118 -

(20  $\mu$ M) to a final volume of about 50  $\mu$ l in a buffer containing 50 mM acetic acid, 50 mM Mes, 100 mM Tris, pH 5.0 and 0.05 percent (w/v) Thesit. The samples were incubated at ambient room temperature for about 6 hours. At the end of the incubation period, reaction mixtures were quenched by addition of about 200  $\mu$ l of 5 percent (v/v) trifluoroacetic acid (TFA). Quenched reactions were analyzed by HPLC on a 5  $\mu$ m, C18 reversed-phase column (Sephasil Peptide, 4.6 x 100 mm; Amersham Pharmacia Biotech, Piscataway, NJ) using first a linear gradient of 18-31.5 percent acetonitrile in 0.1 percent TFA over 40 minutes, then 31.5-81 percent acetonitrile for 1 minute, then 81 percent acetonitrile for 4 minutes, and then returning to 18 percent acetonitrile for several minutes to reequilibrate the column. The linear 18-31.5 percent acetonitrile gradient was subsequently modified to a concave gradient (curve 7 as defined by the Dionex GP40 pump system, Dionex Corporation, Sunnyvale, CA) from 18-31.5 percent acetonitrile in 15 minutes. Proteolysis of the Sw and WT octapeptides was evaluated using a linear gradient of 27-37.8 percent acetonitrile over 24 minutes on the same column.

Both product and substrate were monitored by absorbance at 360 nm. Products were typically identified by retention time comparison with appropriate substrate peptides run under identical

- 119 -

conditions. The rate of product formation or the amount of product formed was determined by comparison of the area under the product peak to a standard curve of authentic peptide. The reference product peptides were each labeled with dinitrophenol at the amino terminus, and had the sequences:

EVNL (SEQ ID NO:24)

EVKV (SEQ ID NO:25)

EVKM (SEQ ID NO:26)

The concentration of standard was determined from the extinction coefficient of DNP ( $\epsilon_{280} = 16 \text{ mM}^{-1}\text{cm}^{-1}$ ). The HPLC assay was linear and reproducible from approximately 25 to about 300 pmol of product.

The results are shown in Figure 11 for each substrate. As can be seen, the Swedish substrate (Sw) is clearly preferred by the enzyme over the wild type (Wt) substrate which is preferred over the methionine to valine mutant (MV) substrate. These results indicate that purified beta-secretase-Fc has the same substrate specificity that has previously been described for beta-secretase from intact cells (Citron et al., *Neuron* 14:661-670 91995!).

While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent

5

- 120 -

variations which come within the scope of the invention  
as claimed.

10

15

20

25

30

35

40

45

50

55

## Claims

5

10

15

20

25

30

35

40

45

50

55

- 121 -

WHAT IS CLAIMED IS:

1. An isolated nucleic acid molecule selected from the group consisting of:
  - a) the nucleic acid molecule as set forth in any of SEQ ID NOS: 1, 2, and 3;
  - b) a nucleic acid molecule encoding the polypeptide of any of SEQ ID NOS: 4, 5, and 6;
  - c) biologically active fragments of SEQ ID NO:4;
  - d) an allelic variant or splice variant of any of (a) or (b);
  - e) a nucleic acid molecule of the DNA vector insert in ATCC Deposit No. 207158;
  - f) a nucleic acid molecule of the DNA vector insert in ATCC Deposit No. 207159;
  - g) a nucleic acid molecule encoding a polypeptide having one to fifty conservative amino acid substitutions as compared with the polypeptide of SEQ ID NO:4, wherein the polypeptide encoded by said nucleic acid molecule is biologically active; and
  - h) a nucleic acid molecule that is the complement of any of (a)-(g) above.
2. A nucleic acid molecule that is SEQ ID NO:1.
3. A nucleic acid molecule encoding the polypeptide of SEQ ID NO:4.
4. A nucleic acid molecule encoding the biologically active polypeptide fragment of claim 13.
5. An expression vector comprising the nucleic acid molecule of Claim 1.

- 122 -

6. A host cell comprising the expression vector of Claim 5.

7. The host cell of Claim 6 which is a eukaryotic cell.

8. The host cell of Claim 6 which is a prokaryotic cell.

9. A process for producing a beta secretase polypeptide comprising culturing the host cell of Claim 6 in suitable culture medium and isolating the polypeptide from the culture.

10. A polypeptide produced by the process of Claim 9.

11. An isolated polypeptide selected from the group consisting of:

- a) the polypeptide of any of SEQ ID NOS: 4, 5, and 6;
- b) a biologically active fragment of any of SEQ ID NOS. 4, 5, 6;
- c) a biologically active polypeptide having one to fifty conservative amino acid changes as compared with the polypeptide of SEQ ID NO:4;
- d) the polypeptide encoded by the DNA vector insert of ATCC Deposit Nos. 207158 and 207159; and
- e) a polypeptide that is an allelic variant or splice variant of (a).

12. An isolated polypeptide encoded by the nucleic acid molecule of Claim 1.

13. An isolated polypeptide having the amino acid sequence of SEQ ID NO:4.

- 123 -

5  
10  
14. An isolated beta secretase polypeptide fragment of SEQ ID NO:4 selected from the group consisting of: amino acids 45-501; amino acids 46-501; amino acids 46-460; amino acids 45-460; amino acids 1-460; amino acids 93-292; amino acids 93-293; amino acids 91-295; amino acids 90-295; amino acids 90-300; amino acids 62-420; amino acids 1-420; amino acids 62-460; amino acids 90-460; amino acids 62-501; amino acids 62-460; amino acids 90-293; amino acids 90-300; amino acids 1-420; amino acids 46-420; amino acids 62-420; amino acids 73-420; amino acids 83-420; amino acids 90-420; amino acids 62-417; amino acids 73-417; amino acids 83-417; amino acids 90-417; amino acids 62-410; amino acids 73-410; amino acids 83-410; amino acids 90-410; amino acids 62-402; amino acids 73-402; amino acids 83-402; and amino acids 90-402.

15  
20  
15. An antibody or fragment thereof specifically binding the polypeptide of Claim 10.

30  
16. The antibody of Claim 15 that is a monoclonal antibody.

35  
25  
17. A composition comprising the polypeptide of Claim 10 and a pharmaceutically acceptable formulation agent.

40  
30  
18. A polypeptide comprising a derivative of the polypeptide of Claim 11.

45  
35  
19. A fusion polypeptide comprising the polypeptide of Claim 11 fused to a heterologous amino acid sequence.

50

55



5

- 124 -

10

20. The fusion polypeptide of Claim 18  
wherein the heterologous amino acid sequence is an IgG  
constant domain or fragment thereof.

15

5 21. A method of modulating levels of beta  
secretase polypeptide in a mammal comprising  
administering to the animal the nucleic acid molecule  
of Claim 1.

20

25

30

35

40

45

50

55

## FIG. 1A

ATGGCCCAAG CCCTGCCCTG GCTCCTGCTG TGGATGGGCG CGGGAGTGCT  
GCCTGCCCAC GGCACCCAGC ACGGCATCCG GCTGCCCCTG CGCAGCGGCC  
TGGGGGGCGC CCCCTGGGG CTGCGGCTGC CCCGGGAGAC CGACGAAGAG  
CCCGAGGAGC CCGGCCGGAG GGGCAGCTTT GTGGAGATGG TGGACAACCT  
GAGGGGCAAG TCGGGGCAGG GCTACTACGT GGAGATGACC GTGGGCAGCC  
CCCCGCAGAC GCTCAACATC CTGGTGGATA CAGGCAGCAG TAACTTTGCA  
GTGGGTGCTG CCCCCACCC CTTCTGTCAT CGCTACTACC AGAGGCAGCT  
GTCCAGCACA TACCGGGACC TCCGAAGGG TGTGTATGTG CCCTACACCC  
AGGGCAAGTG GGAAGGGGAG CTGGGCACCG ACCTGGTAAG CATCCCCCAT  
GGCCCCAACG TCACTGTGCG TGCCAACATT GCTGCCATCA CTGAATCAGA  
CAAGTCTTC ATCAACGGCT CCAACTGGGA AGGCATCTG GGGCTGGCCT  
ATGCTGAGAT TGCCAGGCCT GACGACTCCC TGGAGCCTTT CTTTGA CTCT  
CTGGTAAAGC AGACCCACGT TCCCAACCTC TTCTCCCTGC AGCTTTGTGG  
TGCTGGCTTC CCCCTCAACC AGTCTGAAGT GCTGGCCTCT GTCGGAGGGA  
GCATGATCAT TGGAGGTATC GACCACTCGC TGTACACAGG CAGTCTCTGG  
TATACACCCA TCCGGCGGGA GTGGTATTAT GAGGTGATCA TTGTGCGGGT  
GGAGATCAAT GGACAGGATC TGAAAATGGA CTGCAAGGAG TACA ACTATG  
ACAAGAGCAT TGTGGACAGT GGCACCACCA ACCTTCGTTT GCCCAAGAAA  
GTGTTTGAAG CTGCAGTCAA ATCCATCAAG GCAGCCTCCT CCACGGAGAA  
GTTCCCTGAT GGTTCCTGGC TAGGAGAGCA GCTGGTGTGC TGGCAAGCAG  
GCACCACCCC TTGGAACATT TTCCAGTCA TCTCACTCTA CCTAATGGGT  
GAGGTTACCA ACCAGTCCTT CCGCATCACC ATCCTCCGC AGCAATACCT  
GCGGCCAGTG GAAGATGTGG CCACGTCCCA AGACGACTGT TACAAGTTTG

## FIG. 1B

CCATCTCACA GTCATCCACG GGCACGTGTTA TGGGAGCTGT TATCATGGAG  
GGCTTCTACG TTGTCTTTGA TCGGGCCCGA AAACGAATTG GCTTTGCTGT  
CAGCGCTTGC CATGTGCACG ATGAGTTCAG GACGGCAGCG GTGGAAGGCC  
CTTTTGTCAC CTTGGACATG GAAGACTGTG GCTACAACAT TCCACAGACA  
GATGAGTCAA CCCTCATGAC CATAGCCTAT GTCATGGCTG CCATCTGCGC  
CCTCTTCATG CTGCCACTCT GCCTCATGGT GTGTCAGTGG CGCTGCCTCC  
GCTGCCTGCG CCAGCAGCAT GATGACTTTG CTGATGACAT CTCCCTGCTG  
AAG

## FIG. 2A

ATGGCCCCAG CGCTGCACTG GCTCCTGCTA TGGGTGGGCT CGGGAATGCT  
GCC TGCC CAG GGAACCCATC TCGGCATCCG GCTGCCCCCTT CGCAGCGGCC  
TGGCAGGGCC ACCCCTGGGC CTGAGGCTGC CCCGGGAGAC CGACGAGGAA  
TCGGAGGAGC CTGGCCGGAG AGGCAGCTTT GTGGAGATGG TGGACAACCT  
GAGGGGAAAG TCCGGCCAGG GCTACTATGT GGAGATGACC GTAGGCAGCC  
CCCCACAGAC GCTCAACATC CTGGTGGACA CGGGCAGTAG TAACTTTGCA  
GTGGGGGCTG CCCCACACCC TTTCTGCAT CGCTACTACC AGAGGCAGCT  
GTCCAGCACA TATCGAGACC TCCGAAAGGG TGTGTATGTG CCCTACACCC  
AGGGCAAGTG GGAGGGGGAA CTGGGCACCG ACCTGGTGAG CATCCCTCAT  
GGCCCCAAGC TCACTGTGCG TGCCAAACATT GCTGCCATCA CTGAATCGGA  
CAAGTTCTTC ATCAATGGTT CCAACTGGGA GGGCATCCTA GGGCTGGCCT  
ATGCTGAGAT TGCCAGGCCC GACGACTCTT TGGAGCCCTT CTTTGACTCC  
CTGGTGAAGC AGACCCACAT TCCCAACATC TTTTCCCTGC AGCTCTGTGG  
CGCTGGCTTC CCCCTCAACC AGACCGAGGC ACTGGCCTCG GTGGGAGGGA  
GCATGATCAT TGGTGGTATC GACCACTCGC TATACACGGG CAGTCTCTGG  
TACACACCCA TCCGGCGGGA GTGGTATTAT GAAGTGATCA TTGTACGTGT  
GGAAATCAAT GGTCAAGATC TCAAGATGGA CTGCAAGGAG TACAAC TACG  
ACAAGAGCAT TGTGGACAGT GGGACCACCA ACCTTCGCTT GCCCAAGAAA  
GTATTTGAAG CTGCCGTCAA GTCCATCAAG GCAGCCTCCT CGACGGAGAA  
GTTCCCGGAT GGCTTTTGGC TAGGGGAGCA GCTGGTGTGC TGGCAAGCAG  
GCACGACCCC TTGGAACATT TTCCAGTCA TTCACTTTA CCTCATGGGT  
GAAGTCACCA ATCAGTCCTT CCGCATCACC ATCCTTCCTC AGCAATACCT  
ACGGCCGGTG GAGGACGTGG CCACGTCCCA AGACGACTGT TACAAGTTCC  
CTGTCTCACA GTCATCCACG GGCAGTGTTA TGGGAGCCGT CATCATGGAA

## FIG. 2B

GGTTTCTATG TCGTCTTCGA TCGAGCCCGA AAGCGAATTG GCTTTGCTGT  
CAGCGCTTGC CATGTGCACG ATGAGTTCAG GACGGCGGCA GTGGAAGGTC  
CGTTTGTTAC GGCAGACATG GAAGACTGTG GCTACAACAT TCCCCAGACA  
GATGAGTCAA CACTTATGAC CATAGCCTAT GTCATGGCGG CCATCTGCGC  
CCTCTTCATG TTGCCACTCT GCCTCATGGT ATGTCAGTGG CGCTGCCTGC  
GTTGCCTGCG CCACCAGCAC GATGACTTTG CTGATGACAT CTCCTGCTC  
AAG

## FIG. 3A

ATGGCCCCGG CGCTGCGCTG GCTCCTGCTA TGGGTGGGCT CGGGAATGCT  
GCCTGCCCAG GGAACCCATC TCGGTATCCG ACTGCCCCCTT CGCAGCGGCC  
TGGCAGGGCC ACCCTGGGCG CTGAGGCTGC CCCGGGAGAC GGACGAGGAA  
CCTGAGGAGC CTGGCCGGAG AGGCAGCTTT GTGGAGATGG TGGACAACTT  
GAGGGGAAAAG TCCGGCCAGG GCTACTATGT GGAGATGACC GTGGGCAGCC  
CCCCACAGAC GCTCAACATC CTGGTGGACA CGGGCAGTAG TAATTTTGCA  
GTGGGGGCTG CCCCACACCC TTTCCTGCAT CGATACTACC AAAGGCAGCT  
GTCCAGTACA TACCGAGACC TCCGAAAGTC TGTGTATGTG CCCTACACCC  
AGGGCAAGTG GGAGGGGGAA CTGGGCACTG ACCTGGTGAG CATCCCTCAT  
GGCCCCAACG TCACTGTGCG TGCCAACATT GCTGCCATCA CTGAATCGGA  
CAAGTTCTTC ATCAATGGTT CCAACTGGGA GGGCATCCTA GGGCTGGCCT  
ATGCTGAGAT TGCCAGGCCT GACGACTCCT TGGAGCCCTT TTTTGACTCC  
CTGGTGAAGC AGACCCACAT TCCGAACATC TTTCCCTGC AGCTCTGTGG  
CGCTGCCTTC CCCCTCAACC AGACTGAGGC ACTGGCCTCG GTGGGAGGGA  
GCATGATCAT TGGTGGTATC GACCATTCCC TATACACTGG CAGTCTCTGG  
TACACACCCA TCCGGCGGGA GTGGTATTAT GAAGTGATCA TTGTACGTGT  
AGAAATCAAT GGTCAAGATC TGAAAATGGA CTGCAAGGAG TACAACTATG  
ACAAGAGCAT CGTGGACAGT GGCACCAACA ACCTTCGTTT GCCCAAGAAA  
GTATTTGAAG CTGCAGTCAA GTCCATCAAG GCAGCCTCCT CGACGGAGAA  
GTTCCCGGAT GGCTTTTGGC TAGGGGAGCA GCTGGTGTGC TGGCAAGCAG  
GCACGACCCC TTGGAACATT TTCCAGTCA TTCACTTTA CCTCATGGGT  
GAAGTCACCA ATCAGTCCTT CCGCATCACC ATCCTTCCTC AGCAATACCT  
ACGGCCAGTG GAAGATCTGG CCACGTCCCA AGACGACTGT TACAAGTTCG

## FIG. 3B

CCGTCTCACA GTCATCCACA GGCACCGTTA TGGGAGCGGT CATCATGGAA  
GGCTTCTATG TGGTCTTTGA TCGAGCCCGA AAGCGAATTG GCTTTGCTGT  
CAGCGCTTGC CATGTGCACG ATGAGTTCAG GACGGCGGCA GTGGAAGGTC  
CGTTTGTCAC GGCAGACATG GAAGACTGTG GCTACAACAT TCCACAGACA  
GATGAGTCAA CACTTATGAC CATAGCCTAT GTCATGGCTG CCATCTGCGC  
CCTCTTCATG TTGCCACTCT GCCTCATGGT ATGTCAGTGG CGCTGCCTAC  
GCTGCCTGCG CCATCAGCAT GATGACTTTG CTGATGACAT CTCCTGCTG  
AAA

## FIG. 4

MAQALPWLLL WMCAGVLPAA GTQHGIRLPL RSGLGGAPLG LRLPRETDEE  
PEEPGRGGSF VEMVDNLRGK SGQGYVEMT VGSPPTLNI LVDTGSSNFA  
VGAAPHPFLH RYYQRQLSST YRDLRKGVYV PYTQGWEGE LGTDLVSIPH  
GPNVTVRANI AAITESDKFF INGSNWEGIL GLAYAEIARP DDSLEPFFDS  
LVKQTHVPNL FSLQLCGAGF PLNQSEVLAS VGGSMIIGGI DHSLYTGSIW  
YTPIRREWYY EVIIVRVEIN GQDLKMDCKE YNYDKSIVDS GTTNLRLPKK  
VFEAAVKSIK AASSTEKFPD GFWLGEQLVC WQAGTTPWNI FPVISLYLMG  
EVTNQSFRT ILPQQYLRPV EDVATSQDDC YKFAISQSST GTVMGAVIME  
GFYVVFDRAR KRIGFAVSAC HVHDEFRTAA VEGPFVTLDL EDCGYNIPQT  
DESTLMTIAY VMAAICALFM LPLCLMVCQW RCLRCLRQQH DDFADDISLL  
K




## FIG. 5

MAPALHWLLL WVGSGMLPAQ GTHLGIRLPL RSGLAGPPLG LRLPRETDEE  
SEEPGRRSF VEMVDNLRGK SGQGYVEMT VGSPPTLNI LVDTGSSNFA  
VGAAPHPFLH RYYQRQLSST YRDLRKGYYV PYTQKWEGE LGTDLVSIPH  
GPNVTVRANI AAITESDKFF INGSNWEGIL GLAYAEIARP DDSLEPPFDS  
LVKQTHIPNI FSLQLCGAGF PLNQTEALAS VGGSMIIGGI DHSLYTGSLW  
YTPIRREWYY EVIIVRVEIN GQDLKMDCKE YNYDKSIVDS GTTNLRLPKK  
VFEEAVKSIK AASSTEKFPD GFWLGEQLVC WQAGTPWNI FPVISLYLMG  
EVTNQSFRT ILPQQYLRPV EDVATSQDDC YKFAVSQSST GTVMGAVIME  
GFYVVFDRAR KRIGFAVSAC HVHDEFRTAA VEGPFVTADM EDCGYNIPQT  
DESTLMTIAY VMAAICALFM LPLCLMVCQW RCLRCLRHHQ DDFADDISLL  
K

## FIG. 6

MAPALRWLLL WVGSGMLPAQ GTHLGIRLPL RSGLAGPPLG LRLPRETDEE  
PEEPGRRGSF VEMVDNLRGK SGQGYVEMT VGSPQTLNI LVDTGSSNFA  
VGAAPHFLH RYYQRQLSST YRDLRKSIVV PYTQGWEGE LGTDLVSIPH  
GPNVTVRANI AAITESDKFF INGSNWEGIL GLAYAEIARP DDSLEPPFDS  
LVKQTHIPNI FSLQLCGAGF PLNQTEALAS VGGSMIIGGI DHSLYTGSW  
YTPIRREWYY EVIIVRVEIN GQDLKMDCKE YNYDKSIVDS GTTNLRLPKK  
VFEAAVKSIX AASSTEKFPD GFWLGEQLVC WQAGTTPWNI FPVISLYLMG  
EVTNQSFRIIT ILPQQYLRPV EDVATSQDDC YKFAVSQSST GTVMGAVIME  
GFYVVFDRAR KRIGFAVSAC HVHDEFRTAA VEGPFVTADM EDCGYNIPQT  
DESTLMTIAY VMAAICALFM LPLCLMVCQW RCLRCLRHQH DDFADDISLL

K






FIG. 9A




FIG. 9B




FIG. 10




FIG. 11





FIG. 12





# INTERNATIONAL SEARCH REPORT

national Application No  
PCT/US 00/07755

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C12N15/52 C12N9/64 C07K16/40 A61K38/48 C12N15/62 A01K67/027 G01N33/566 A61K48/00		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C12N C07K A61K A01K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) STRAND, BIOSIS, EMBASE, MEDLINE, CAB Data, WPI Data, EPO-Internal, PAJ		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
E	WO 00 17369 A (PARODI LUIS A ;BIENKOWSKI MICHAEL JEROME (US); HEINRIKSON ROBERT L) 30 March 2000 (2000-03-30) the whole document	1-21
P,X	SINHA S ET AL: "PURIFICATION AND CLONING OF AMYLOID PRECURSOR PROTEIN BETA-SECRETASE FROM HUMAN BRAIN" NATURE,GB,MACMILLAN JOURNALS LTD. LONDON, vol. 402, no. 6761, 2 December 1999 (1999-12-02), pages 537-540, XP000881765 ISSN: 0028-0836 the whole document	1-21
--- -/--		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of box C.		
<input checked="" type="checkbox"/> Patent family members are listed in annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "Z" document member of the same patent family		
Date of the actual completion of the international search 10 July 2000		Date of mailing of the international search report 24/07/2000
Name and mailing address of the ISA European Patent Office, P.O. 5618 Patentaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Te. 31 651 epo nl, Fax: (+31-70) 340-3010		Authorized officer Hix, R

Form PCT/ISA/210 (second sheet) July 1992

# INTERNATIONAL SEARCH REPORT

national Application No

PCT/US 00/07755

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant part	Relevant to claim No.
P,X	R. YAN ET AL.: "Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity." NATURE, vol. 402, no. 6761, 2 December 1999 (1999-12-02), pages 533-537, XP002141788 the whole document	1-21
P,X	I. HUSSAIN ET AL.: "Identification of a novel aspartic protease (Asp 2) as Beta-secretase." MOLECULAR AND CELLULAR NEUROSCIENCE, vol. 14, no. 6, December 1999 (1999-12), pages 419-427, XP000921315 the whole document	1-21
P,X	R. VASSAR ET AL.: "Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE." SCIENCE, vol. 286, no. 5440, 22 October 1999 (1999-10-22), pages 735-741, XP000914811 the whole document	1-21
X	EP 0 855 444 A (SMITHKLINE BEECHAM PLC ;SMITHKLINE BEECHAM CORP (US)) 29 July 1998 (1998-07-29) cited in the application the whole document	1-21
Y	US 5 744 346 A (KEIM PAMELA S ET AL) 28 April 1998 (1998-04-28) cited in the application the whole document	1-21
Y	WO 98 26059 A (KEIM PAMELA S ;SINHA SUKANTO (US); ANDERSON JOHN P (US); ATHENA NE) 18 June 1998 (1998-06-18) the whole document	1-21
P,Y	WO 99 64587 A (BERTRAND PHILIPPE ;MOUTAOUAKIL MOHAMED (FR); MUNOZ GIMENEZ NOELI ( ) 16 December 1999 (1999-12-16) the whole document	1-21
	--- -/--	

# INTERNATIONAL SEARCH REPORT

national Application No  
PCT/US 00/07755

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim no.
A	G. HUBER ET AL.: "cDNA cloning and molecular characterization of human brain metalloprotease MP100: A beta-secretase candidate?" JOURNAL OF NEUROCHEMISTRY, vol. 72, no. 3, March 1999 (1999-03), pages 1215-1223, XP000921316 the whole document	
A	A. THOMPSON ET AL.: "Expression and characterization of human beta-secretase candidates metalloendopeptidase MP78 and cathepsin D in beta-APP- overexpressing cells" MOLECULAR BRAIN RESEARCH, vol. 48, no. 2, 1997, pages 206-214, XP000923393 the whole document	
A	N. CHEVALLIER ET AL.: "Cathepsin D displays in vitro beta-secretase-like specificity." BRAIN RESEARCH, vol. 750, 1997, pages 11-19, XP000921314 the whole document	
A	EP 0 848 062 A (SMITHKLINE BEECHAM PLC ; SMITHKLINE BEECHAM CORP (US)) 17 June 1998 (1998-06-17) cited in the application the whole document	

# INTERNATIONAL SEARCH REPORT

Information on patent family members

National Application No

PCT/US 00/07755

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0017369 A	30-03-2000	AU 6141899 A	10-04-2000
EP 0855444 A	29-07-1998	CA 2221686 A	28-07-1998
		JP 10327875 A	15-12-1998
		JP 2000060579 A	29-02-2000
US 5744346 A	28-04-1998	AU 6383396 A	30-12-1996
		EP 0871720 A	21-10-1998
		JP 11507538 T	06-07-1999
		WO 9640885 A	19-12-1996
		US 5942400 A	24-08-1999
WO 9826059 A	18-06-1998	AU 1684097 A	03-07-1998
WO 9964587 A	16-12-1999	FR 2779444 A	10-12-1999
		AU 4045599 A	30-12-1999
EP 0848062 A	17-06-1998	JP 11069981 A	16-03-1999
		US 6025180 A	15-02-2000

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

**BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☒ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☒ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**